

Anti-SULT2A1 Picoband Antibody

Catalog # ABO12507

Product Information

Application	WB, IHC-P
Primary Accession	Q06520
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Bile salt sulfotransferase(SULT2A1) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	6822
Other Names	Bile salt sulfotransferase, 2.8.2.14, Dehydroepiandrosterone sulfotransferase, DHEA-ST, Hydroxysteroid Sulfotransferase, HST, ST2, ST2A3, Sulfotransferase 2A1, ST2A1, SULT2A1, HST, STD
Calculated MW	33780
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 μ g/ml, Human, Mouse, Rat
Subcellular Localization	Cytoplasm.
Tissue Specificity	Liver, adrenal and at lower level in the kidney. Is present in human fetus in higher level in the adrenal than the liver and the kidney.
Source	Eukaryota
Protein Name	Bile salt sulfotransferase
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human SULT2A1 (253-285aa DWKNHFTVAQAEDFDKLFQEKMAADLPRELPWE), different from the related mouse sequence by seven amino acids, and from the related rat sequence by eight amino acids.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins
Storage	At -20° C for one year. After r° Constitution, at 4° C for one month. It° Can also be aliquotted and stored frozen at -20° C for a longer time.Avoid repeated freezing and thawing.

Protein Information



Name	SULT2A1
Synonyms	HST, STD
Function	Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the sulfonation of steroids and bile acids in the liver and adrenal glands (PubMed: 14573603 , PubMed: 18042734 , PubMed: 19589875 , PubMed: 20102295 , PubMed: 21187059 , PubMed: 2268288 , PubMed: 29671343 , PubMed: 7678732 , PubMed: 7854148). Mediates the sulfation of a wide range of steroids and sterols, including pregnenolone, androsterone, DHEA, bile acids, cholesterol and as well many xenobiotics that contain alcohol and phenol functional groups (PubMed: 14573603 , PubMed: 18042734 , PubMed: 19589875 , PubMed: 20102295 , PubMed: 21187059 , PubMed: 2268288 , PubMed: 29671343 , PubMed: 7678732 , PubMed: 7854148). Sulfonation increases the water solubility of most compounds, and therefore their renal excretion, but it can also result in bioactivation to form active metabolites. Plays an important role in maintaining steroid and lipid homeostasis (PubMed: 14573603 , PubMed: 19589875 , PubMed: 21187059). Plays a key role in bile acid metabolism, mediating formation of 3-sulfated bile acids (PubMed: 2268288 , PubMed: 20102295). In addition, catalyzes the metabolic activation of potent carcinogenic polycyclic arylmethanols (By similarity).
Cellular Location	Cytoplasm.
Tissue Location	Liver, adrenal and at lower level in the kidney. Is present in human fetus in higher level in the adrenal than the liver and the kidney

Background


Bile salt sulfotransferase, also known as hydroxysteroid sulfotransferase (HST) or sulfotransferase 2A1 (ST2A1), is an enzyme that in humans is encoded by the SULT2A1 gene. It is mapped to 19q13.3. This gene encodes a member of the sulfotransferase family. Sulfotransferases aid in the metabolism of drugs and endogenous compounds by converting these substances into more hydrophilic water-soluble sulfate conjugates that can be easily excreted. This protein catalyzes the sulfation of steroids and bile acids in the liver and adrenal glands, and may have a role in the inherited adrenal androgen excess in women with polycystic ovary syndrome.

Images


Anti- SULT2A1 Picoband antibody, ABO12507, Western blottingAll lanes: Anti SULT2A1 (ABO12507) at 0.5ug/mlLane 1: Rat Liver Tissue Lysate at 50ugLane 2: Rat Kidney Tissue Lysate at 50ugLane 3: Mouse Liver Tissue Lysate at 50ugLane 4: Mouse Kidney Tissue Lysate at 50ugLane 5: HELA Whole Cell Lysate at 40ugLane 6: SW620 Whole Cell Lysate at 40ugPredicted bind size: 34KDObserved bind size: 34KD



Anti- SULT2A1 Picoband antibody, ABO12507,
IHC(P)IHC(P): Mouse Liver Tissue



Anti- SULT2A1 Picoband antibody, ABO12507,
IHC(P)IHC(P): Rat Liver Tissue



Anti- SULT2A1 Picoband antibody, ABO12507,
IHC(P)IHC(P): Human Renal Cancer Tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.