

Anti-BCAR3 Picoband Antibody

Catalog # ABO12532

Product Information

Application	WB, IHC-P
Primary Accession	O75815
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Breast cancer anti-estrogen resistance protein 3(BCAR3) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	8412
Other Names	Breast cancer anti-estrogen resistance protein 3, Novel SH2-containing protein 2, SH2 domain-containing protein 3B, BCAR3, NSP2, SH2D3B
Calculated MW	92566
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5 µg/ml, Human
Tissue Specificity	Ubiquitously expressed. Found in several cancer cell lines, but not in nonmalignant breast tissue. .
Source	Eukaryota
Protein Name	Breast cancer anti-estrogen resistance protein 3
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human BCAR3 (791-825aa KGAQVNQTERYEKFNQILTALSRKLEPPPVKQAEEL), different from the related mouse sequence by five amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also

be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

Protein Information

Name	BCAR3
Synonyms	NSP2, SH2D3B
Function	Acts as an adapter protein downstream of several growth factor receptors to promote cell proliferation, migration, and redistribution of actin fibers (PubMed: 24216110). Specifically involved in INS/insulin signaling pathway by mediating MAPK1/ERK2-MAPK3/ERK1 activation and DNA synthesis (PubMed: 24216110). Promotes insulin- mediated membrane ruffling (By similarity). In response to vasoconstrictor peptide EDN1, involved in the activation of RAP1 downstream of PTK2B via interaction with phosphorylated BCAR1 (PubMed: 19086031). Inhibits cell migration and invasion via regulation of TGFB-mediated matrix digestion, actin filament rearrangement, and inhibition of invadopodia activity (By similarity). May inhibit TGFB- SMAD signaling, via facilitating BCAR1 and SMAD2 and/or SMAD3 interaction (By similarity). Regulates EGF-induced DNA synthesis (PubMed: 18722344). Required for the maintenance of ocular lens morphology and structural integrity, potentially via regulation of focal adhesion complex signaling (By similarity). Acts upstream of PTPRA to regulate the localization of BCAR1 and PTPRA to focal adhesions, via regulation of SRC-mediated phosphorylation of PTPRA (By similarity). Positively regulates integrin-induced tyrosine phosphorylation of BCAR1 (By similarity). Acts as a guanine nucleotide exchange factor (GEF) for small GTPases RALA, RAP1A and RRAS (By similarity). However, in a contrasting study, lacks GEF activity towards RAP1 (PubMed: 22081014).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:Q9QZK2}. Cell junction, focal adhesion {ECO:0000250 UniProtKB:Q9QZK2} Note=Localization to focal adhesions depends on interaction with PTPRA {ECO:0000250 UniProtKB:Q9QZK2}
Tissue Location	Ubiquitously expressed. Found in several cancer cell lines, but not in nonmalignant breast tissue

Background


Breast cancer anti-estrogen resistance protein 3 is a protein that in humans is encoded by the BCAR3 gene. Breast tumors are initially dependent on estrogens for growth and progression and can be inhibited by anti-estrogens such as tamoxifen. However, breast cancers progress to become anti-estrogen resistant. Breast cancer anti-estrogen resistance gene 3 was identified in the search for genes involved in the development of estrogen resistance. The gene encodes a component of intracellular signal transduction that causes estrogen-independent proliferation in human breast cancer cells. The protein contains a putative src homology 2 (SH2) domain, a hall mark of cellular tyrosine kinase signaling molecules, and is partly homologous to the cell division cycle protein CDC48. Multiple transcript variants encoding different isoforms have been found for this gene.

Images


Western blot analysis of BCAR3 expression in HEPG2 whole cell lysates (lane 1). BCAR3 at 93KD was detected using rabbit anti- BCAR3 Antigen Affinity purified polyclonal antibody(Catalog # ABO12532) at 0.5 ??g/mL.




The blot was developed using chemiluminescence (ECL) method .



BCAR3 was detected in paraffin-embedded sections of mouse lymphaden tissues using rabbit anti- BCAR3 Antigen Affinity purified polyclonal antibody (Catalog # ABO12532) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .



BCAR3 was detected in paraffin-embedded sections of rat spleen tissues using rabbit anti- BCAR3 Antigen Affinity purified polyclonal antibody (Catalog # ABO12532) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .



BCAR3 was detected in paraffin-embedded sections of human tonsil tissues using rabbit anti- BCAR3 Antigen Affinity purified polyclonal antibody (Catalog # ABO12532) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.