

Anti-SPTLC1 Picoband Antibody

Catalog # ABO12580

Product Information

Application	WB, IHC-P
Primary Accession	O15269
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Serine palmitoyltransferase 1(SPTLC1) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	10558
Other Names	Serine palmitoyltransferase 1, 2.3.1.50, Long chain base biosynthesis protein 1, LCB 1, Serine-palmitoyl-CoA transferase 1, SPT 1, SPT1, SPTLC1, LCB1
Calculated MW	52744
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, By Heat Western blot, 0.1-0.5 μ g/ml, Human, Mouse, Rat
Subcellular Localization	Endoplasmic reticulum membrane ; Single-pass membrane protein .
Tissue Specificity	Widely expressed. Not detected in small intestine. .
Source	Eukaryota
Protein Name	Serine palmitoyltransferase 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ N.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human SPTLC1 (444-473aa IRVWTVETQTEELERAASTIKEVAQAVLL), different from the related mouse sequence by three amino acids.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins.

Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
----------------	---

Protein Information


Name	SPTLC1
Synonyms	LCB1
Function	Component of the serine palmitoyltransferase multisubunit enzyme (SPT) that catalyzes the initial and rate-limiting step in sphingolipid biosynthesis by condensing L-serine and activated acyl-CoA (most commonly palmitoyl-CoA) to form long-chain bases. The SPT complex is also composed of SPTLC2 or SPTLC3 and SPTSSA or SPTSSB. Within this complex, the heterodimer with SPTLC2 or SPTLC3 forms the catalytic core (PubMed: 19416851 , PubMed: 33558762 , PubMed: 36170811). The composition of the serine palmitoyltransferase (SPT) complex determines the substrate preference (PubMed: 19416851 , PubMed: 33558762). The SPTLC1-SPTLC2-SPTSSA complex shows a strong preference for C16-CoA substrate, while the SPTLC1-SPTLC3-SPTSSA isozyme uses both C14-CoA and C16-CoA as substrates, with a slight preference for C14-CoA (PubMed: 19416851 , PubMed: 19648650). The SPTLC1-SPTLC2-SPTSSB complex shows a strong preference for C18-CoA substrate, while the SPTLC1-SPTLC3-SPTSSB isozyme displays an ability to use a broader range of acyl-CoAs, without apparent preference (PubMed: 19416851 , PubMed: 19648650 , PubMed: 33558761 , PubMed: 33558762). Required for adipocyte cell viability and metabolic homeostasis (By similarity).
Cellular Location	Endoplasmic reticulum membrane; Single-pass membrane protein {ECO:0000250 UniProtKB:O35704}
Tissue Location	Widely expressed. Not detected in small intestine.

Background

SPTLC1(Serine palmitoyltransferase, long chain base subunit 1), also known as SPT1, LCB1, is a protein which in humans is encoded by the SPTLC1 gene. Dawkins et al. (2001) noted that the SPTLC1 gene maps to chromosome 9q22.1-q22.3. Serine palmitoyltransferase, which consists of two different subunits, is the initial enzyme in sphingolipid biosynthesis. It converts L-serine and palmitoyl CoA to 3-oxosphinganine with pyridoxal 5'-phosphate as a cofactor. The product of this gene is the long chain base subunit 1 of serine palmitoyltransferase. Mutations in this gene were identified in patients with hereditary sensory neuropathy type 1. Alternatively spliced variants encoding different isoforms have been identified.

Images

Western blot analysis of SPTLC1 expression in rat brain extract (lane 1), mouse brain extract (lane 2) and MCF-7 whole cell lysates (lane 3). SPTLC1 at 53KD was detected using rabbit anti- SPTLC1 Antigen Affinity purified polyclonal antibody (Catalog # ABO12580) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .



SPTLC1 was detected in paraffin-embedded sections of human intestinal cancer tissues using rabbit anti- SPTLC1 Antigen Affinity purified polyclonal antibody (Catalog # ABO12580) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.