

Anti-Factor B Picoband Antibody

Catalog # ABO12619

Product Information

Application	WB, IHC-P
Primary Accession	P00751
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Complement factor B(CFB) detection. Tested with WB, IHC-P in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	629
Other Names	Complement factor B, 3.4.21.47, C3/C5 convertase, Glycine-rich beta glycoprotein, GBG, PBF2, Properdin factor B, Complement factor B Ba fragment, Complement factor B Bb fragment, CFB, BF, BFD
Calculated MW	85533
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat Western blot, 0.1-0.5 µg/ml, Human
Subcellular Localization	Secreted.
Source	Eukaryota
Protein Name	Complement factor B
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E. coli-derived human Factor B recombinant protein (Position: E518-L764). Human Factor B shares 84.2% amino acid (aa) sequence identity with mouse Factor B.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also

be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

Protein Information

Name	CFB (HGNC:1037)
Synonyms	BF, BFD
Function	Precursor of the catalytic component of the C3 and C5 convertase complexes of the alternative pathway of the complement system, a cascade of proteins that leads to phagocytosis and breakdown of pathogens and signaling that strengthens the adaptive immune system (PubMed: 3638964 , PubMed: 624565 , PubMed: 6554279 , PubMed: 6919543 , PubMed: 9748277). The alternative complement pathway acts as an amplification loop that enhances other complement pathways (classical, lectin and GZMK) by promoting formation of additional C3 and C5 convertases (PubMed: 3638964 , PubMed: 624565 , PubMed: 6554279 , PubMed: 6919543 , PubMed: 9748277). CFB is cleaved and activated by CFD to generate Ba and Bb chains; Bb chain constituting the catalytic component of the C3 and C5 convertases (PubMed: 6769474 , PubMed: 9748277).
Cellular Location	Secreted.

Background


Complement factor B is a protein that in humans is encoded by the CFB gene. This gene encodes complement factor B, a component of the alternative pathway of complement activation. Factor B circulates in the blood as a single chain polypeptide. Upon activation of the alternative pathway, it is cleaved by complement factor D yielding the noncatalytic chain Ba and the catalytic subunit Bb. The active subunit Bb is a serine protease which associates with C3b to form the alternative pathway C3 convertase. Bb is involved in the proliferation of preactivated B lymphocytes, while Ba inhibits their proliferation. This gene localizes to the major histocompatibility complex (MHC) class III region on chromosome 6. This cluster includes several genes involved in regulation of the immune reaction. Polymorphisms in this gene are associated with a reduced risk of age-related macular degeneration. The polyadenylation site of this gene is 421 bp from the 5' end of the gene for complement component 2.

Images



Western blot analysis of Factor B expression in A549 whole cell lysates (lane 1), 293T whole cell lysates (lane 2) and HELA whole cell lysates (lane 3). Factor B at 86KD was detected using rabbit anti- Factor B Antigen Affinity purified polyclonal antibody (Catalog # ABO12619) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .

Factor B was detected in paraffin-embedded sections of human lung cancer tissues using rabbit anti- Factor B Antigen Affinity purified polyclonal antibody (Catalog # ABO12619) at 1 µg/mL. The immunohistochemical



section was developed using SABC method .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.