

Anti-WDR1 Picoband Antibody

Catalog # ABO12647

Product Information

Application	WB
Primary Accession	O75083
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for WD repeat-containing protein 1(WDR1) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	9948
Other Names	WD repeat-containing protein 1, Actin-interacting protein 1, AIP1, NORI-1, WDR1
Calculated MW	66194
Application Details	Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat
Subcellular Localization	Cytoplasm, cytoskeleton . Cell projection, podosome . Cell junction .
Source	Eukaryota
Protein Name	WD repeat-containing protein 1
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human WDR1 (552-579aa MDMMVYVWTLSDPETRVKIQDAHRLHHV), different from the related mouse and rat sequences by one amino acid.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.


Protein Information

Name	WDR1
Function	Induces disassembly of actin filaments in conjunction with ADF/cofilin family proteins (PubMed: 15629458 , PubMed: 27557945 , PubMed: 29751004). Enhances cofilin-mediated actin severing (By similarity). Involved in cytokinesis. Involved in chemotactic cell migration by restricting lamellipodial membrane protrusions (PubMed: 18494608). Involved in myocardium sarcomere organization. Required for cardiomyocyte growth and maintenance (By similarity). Involved in megakaryocyte maturation and platelet shedding. Required for the establishment of planar cell polarity (PCP) during follicular epithelium development and for cell shape changes during PCP; the function seems to implicate cooperation with CFL1 and/or DSTN/ADF. Involved in the generation/maintenance of cortical tension (By similarity). Involved in assembly and maintenance of epithelial apical cell junctions and plays a role in the organization of the perijunctional actomyosin belt (PubMed: 25792565).
Cellular Location	Cytoplasm. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:Q5RKI0}. Cell projection, podosome. Cell junction
Tissue Location	Expressed in peripheral blood mononuclear cells (at protein level).

Background

WD repeat-containing protein 1 is a protein that in humans is encoded by the WDR1 gene. It is mapped to 4p16.1. This gene encodes a protein containing 9 WD repeats. WD repeats are approximately 30- to 40-amino acid domains containing several conserved residues, mostly including a trp-asn at the C-terminal end. WD domains are involved in protein-protein interactions. The encoded protein may help induce the disassembly of actin filaments. Two transcript variants encoding different isoforms have been found for this gene.

Images



Western blot analysis of WDR1 expression in rat liver extract (lane 1), mouse brain extract (lane 2) and A431 whole cell lysates (lane 3). WDR1 at 66KD was detected using rabbit anti-WDR1 Antigen Affinity purified polyclonal antibody (Catalog # ABO12647) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.