

Anti-ca2 Picoband Antibody

Catalog # ABO12674

Product Information

Application	WB, IHC-P
Primary Accession	P00918
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Carbonic anhydrase 2(CA2) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	760
Other Names	Carbonic anhydrase 2, 4.2.1.1, Carbonate dehydratase II, Carbonic anhydrase C, CAC, Carbonic anhydrase II, CA-II, CA2
Calculated MW	29246
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, By Heat Western blot, 0.1-0.5 μ g/ml, Human, Mouse, Rat
Subcellular Localization	Cytoplasm . Cell membrane . Colocalized with SLC26A6 at the surface of the cell membrane in order to form a bicarbonate transport metabolon. Displaced from the cytosolic surface of the cell membrane by PKC in phorbol myristate acetate (PMA)-induced cells.
Source	Eukaryota
Protein Name	Carbonic anhydrase 2
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E.coli-derived human CA2 recombinant protein (Position: S2-K260). Human CA2 shares 81.1% and 80.7% amino acid (aa) sequence identity with mouse and rat CA2, respectively.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins

Storage	At -20 °C for one year. After r° Constitution, at 4 °C for one month. It° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.
----------------	---

Protein Information



Name	CA2
Function	Catalyzes the reversible hydration of carbon dioxide (PubMed: 11327835 , PubMed: 11802772 , PubMed: 11831900 , PubMed: 12056894 , PubMed: 12171926 , PubMed: 1336460 , PubMed: 14736236 , PubMed: 15300855 , PubMed: 15453828 , PubMed: 15667203 , PubMed: 15865431 , PubMed: 16106378 , PubMed: 16214338 , PubMed: 16290146 , PubMed: 16686544 , PubMed: 16759856 , PubMed: 16807956 , PubMed: 17127057 , PubMed: 17251017 , PubMed: 17314045 , PubMed: 17330962 , PubMed: 17346964 , PubMed: 17540563 , PubMed: 17588751 , PubMed: 17705204 , PubMed: 18024029 , PubMed: 18162396 , PubMed: 18266323 , PubMed: 18374572 , PubMed: 18481843 , PubMed: 18618712 , PubMed: 18640037 , PubMed: 18942852 , PubMed: 1909891 , PubMed: 1910042 , PubMed: 19170619 , PubMed: 19186056 , PubMed: 19206230 , PubMed: 19520834 , PubMed: 19778001 , PubMed: 7761440 , PubMed: 7901850 , PubMed: 8218160 , PubMed: 8262987 , PubMed: 8399159 , PubMed: 8451242 , PubMed: 8485129 , PubMed: 8639494 , PubMed: 9265618 , PubMed: 9398308). Can also hydrate cyanamide to urea (PubMed: 10550681 , PubMed: 11015219). Stimulates the chloride-bicarbonate exchange activity of SLC26A6 (PubMed: 15990874). Essential for bone resorption and osteoclast differentiation (PubMed: 15300855). Involved in the regulation of fluid secretion into the anterior chamber of the eye. Contributes to intracellular pH regulation in the duodenal upper villous epithelium during proton-coupled peptide absorption.
Cellular Location	Cytoplasm. Cell membrane. Note=Colocalized with SLC26A6 at the surface of the cell membrane in order to form a bicarbonate transport metabolon. Displaced from the cytosolic surface of the cell membrane by PKC in phorbol myristate acetate (PMA)-induced cells

Background


CA2 is a cytosolic enzyme with the highest activity among all known CAs. The carbonic anhydrases (ACs) form a family of enzymes that catalyze the rapid interconversion of carbon dioxide and water to bicarbonate and protons (or vice versa), a reversible reaction that occurs relatively slowly in the absence of a catalyst. Mutations in the CA2 gene result in the CA II deficiency syndrome, an autosomal recessive disorder that produces osteopetrosis, renal tubular acidosis and cerebral calcification. This gene is mapped to 8q22.

Images

Western blot analysis of CA2 expression in rat ovary extract (lane 1), mouse liver extract (lane 2) and MCF-7 whole cell lysates (lane 3). CA2 at 29KD was detected using rabbit anti- CA2 Antigen Affinity purified polyclonal antibody (Catalog # ABO12674) at 0.5 ??g/mL. The blot was developed using chemiluminescence (ECL) method .



CA2 was detected in paraffin-embedded sections of human gastric cancer tissues using rabbit anti- CA2 Antigen Affinity purified polyclonal antibody (Catalog # ABO12674) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .



CA2 was detected in paraffin-embedded sections of human liver cancer tissues using rabbit anti- CA2 Antigen Affinity purified polyclonal antibody (Catalog # ABO12674) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.