

Anti-Growth Hormone/GH1 Antibody

Catalog # ABO12692

Product Information

Application	WB, IHC-P, E
Primary Accession	P01244
Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Somatotropin(GH1) detection. Tested with WB, IHC-P, ELISA in Rat.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	24391
Other Names	Somatotropin, Growth hormone, Gh1, Gh
Calculated MW	24656
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Rat, By Heat ELISA , 0.1-0.5 μ g/ml, Rat, - Western blot, 0.1-0.5 μ g/ml, Rat
Subcellular Localization	Secreted.
Source	Eukaryota
Protein Name	Somatotropin
Contents	Each vial contains 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ . Carrier free (No BSA) form available in stock. If you want this antibody carrier free please specify Carrier Free" or "No BSA" in your order note. "
Immunogen	E. coli-derived rat GH recombinant protein(Position: F27-F216).
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time.Avoid repeated freezing and thawing.


Protein Information

Name	Gh1
Synonyms	Gh
Function	Plays an important role in growth control. Its major role in stimulating body growth is to stimulate the liver and other tissues to secrete IGF1. It stimulates both the differentiation and proliferation of myoblasts. It also stimulates amino acid uptake and protein synthesis in muscle and other tissues.
Cellular Location	Secreted.

Background

Growth Hormone(GH) is mapped to 17q22-q24. Human growth hormone has a molecular mass of 22,005 and contains 191 amino acid residues with 2 disulfide bridges. Rat GH shares 98% amino acid sequence homology with mouse. It binds two receptor molecules and thereby induces signal transduction through receptor dimerization. At high concentrations, GH acts as an antagonist because of a large difference in affinities at the respective binding sites.

Images



Anti-rat Growth Hormone antibody, ABO12692, Western blotting
Lane 1: Recombinant Rat GH Protein 10ng
Lane 2: Recombinant Rat GH Protein 5ng
Lane 3: Recombinant Rat GH Protein 2

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.