

Anti-Livin Antibody

Catalog # ABO12759

Product Information

Application	WB, E
Primary Accession	Q96CA5
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Baculoviral IAP repeat-containing protein 7(BIRC7) detection. Tested with WB, ELISA in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	79444
Other Names	Baculoviral IAP repeat-containing protein 7, 2.3.2.27, Kidney inhibitor of apoptosis protein, KIAP, Livin, Melanoma inhibitor of apoptosis protein, ML-IAP, RING finger protein 50, RING-type E3 ubiquitin transferase BIRC7, Baculoviral IAP repeat-containing protein 7 30kDa subunit, Truncated livin, p30-Livin, tLivin, BIRC7, KIAP, LIVIN, MLIAP, RNF50
Calculated MW	32798
Application Details	Western blot, 0.1-0.5 μ g/ml, Human, - ELISA , 0.1-0.5 μ g/ml, Human
Subcellular Localization	Nucleus . Cytoplasm . Golgi apparatus . Nuclear, and in a filamentous pattern throughout the cytoplasm. Full-length livin is detected exclusively in the cytoplasm, whereas the truncated form (tLivin) is found in the peri-nuclear region with marked localization to the Golgi apparatus; the accumulation of tLivin in the nucleus shows positive correlation with the increase in apoptosis.
Tissue Specificity	Isoform 1 and isoform 2 are expressed at very low levels or not detectable in most adult tissues. Detected in adult heart, placenta, lung, lymph node, spleen and ovary, and in several carcinoma cell lines. Isoform 2 is detected in fetal kidney, heart and spleen, and at lower levels in adult brain, skeletal muscle and peripheral blood leukocytes.
Source	Eukaryota
Protein Name	Baculoviral IAP repeat-containing protein 7

Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Immunogen	E. coli-derived human Livin recombinant protein (Position: E87-S298). Human Livin shares 69.8% amino acid (aa) sequence identity with mouse Livin.
Purification	Immunogen affinity purified.
Cross Reactivity	No cross reactivity with other proteins
Storage	At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.
Sequence Similarities	Belongs to the IAP family.

Protein Information

Name	BIRC7
Synonyms	KIAP, LIVIN, MLIAP, RNF50
Function	Apoptotic regulator capable of exerting proapoptotic and anti-apoptotic activities and plays crucial roles in apoptosis, cell proliferation, and cell cycle control (PubMed: 11024045 , PubMed: 11084335 , PubMed: 11162435 , PubMed: 16729033 , PubMed: 17294084). Its anti-apoptotic activity is mediated through the inhibition of CASP3, CASP7 and CASP9, as well as by its E3 ubiquitin-protein ligase activity (PubMed: 11024045 , PubMed: 16729033). As it is a weak caspase inhibitor, its anti-apoptotic activity is thought to be due to its ability to ubiquitinate DIABLO/SMAC targeting it for degradation thereby promoting cell survival (PubMed: 16729033). May contribute to caspase inhibition, by blocking the ability of DIABLO/SMAC to disrupt XIAP/BIRC4-caspase interactions (PubMed: 16729033). Protects against apoptosis induced by TNF or by chemical agents such as adriamycin, etoposide or staurosporine (PubMed: 11084335 , PubMed: 11162435 , PubMed: 11865055). Suppression of apoptosis is mediated by activation of MAPK8/JNK1, and possibly also of MAPK9/JNK2 (PubMed: 11865055). This activation depends on TAB1 and MAP3K7/TAK1 (PubMed: 11865055). In vitro, inhibits CASP3 and proteolytic activation of pro-CASP9 (PubMed: 11024045).
Cellular Location	Nucleus. Cytoplasm. Golgi apparatus. Note=Nuclear, and in a filamentous pattern throughout the cytoplasm. Full-length livin is detected exclusively in the cytoplasm, whereas the truncated form (tLivin) is found in the peri-nuclear region with marked localization to the Golgi apparatus; the accumulation of tLivin in the nucleus shows positive correlation with the increase in apoptosis
Tissue Location	Isoform 1 and isoform 2 are expressed at very low levels or not detectable in most adult tissues. Detected in adult heart, placenta, lung, lymph node, spleen and ovary, and in several carcinoma cell lines. Isoform 2 is detected in fetal kidney, heart and spleen, and at lower levels in adult brain, skeletal muscle and peripheral blood leukocytes

Background

Baculoviral IAP repeat-containing protein 7 is a protein that in humans is encoded by the BIRC7 gene. This protein is a member of the family of inhibitor of apoptosis proteins (IAP) and contains a single copy of a baculovirus IAP repeat (BIR) and a RING finger motif. Its gene is mapped to 20q13.3. It has got 280- and 298-amino acids. The protein is highly expressed in melanomas while lowly expressed in some lymphomas, fetal

kidney, fetal liver, testis and thymus. BIRC7 plays a vital role in blocking apoptosis induced.

Images



Anti- Livin antibody, ABO12759, Western blotting
All lanes:
Anti Livin (ABO12759) at 0.5ug/ml
Lane 1: HELA Whole Cell Lysate at 40ug
Lane 2: A431 Whole Cell Lysate at 40ug
Predicted bind size: 33KD
Observed bind size: 33KD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.