

# Anti-Syndecan-1/SDC1 Picoband Antibody

Catalog # ABO12911

## Product Information

---

|                          |                                                                                         |
|--------------------------|-----------------------------------------------------------------------------------------|
| <b>Application</b>       | WB                                                                                      |
| <b>Primary Accession</b> | <a href="#">P18827</a>                                                                  |
| <b>Host</b>              | Rabbit                                                                                  |
| <b>Reactivity</b>        | Human                                                                                   |
| <b>Clonality</b>         | Polyclonal                                                                              |
| <b>Format</b>            | Lyophilized                                                                             |
| <b>Description</b>       | Rabbit IgG polyclonal antibody for Syndecan-1(SDC1) detection. Tested with WB in Human. |

**Reconstitution** Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

## Additional Information

---

|                                 |                                                                                                                                                                                                                  |
|---------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Gene ID</b>                  | 6382                                                                                                                                                                                                             |
| <b>Other Names</b>              | Syndecan-1, SYND1, CD138, SDC1, SDC                                                                                                                                                                              |
| <b>Calculated MW</b>            | 32462                                                                                                                                                                                                            |
| <b>Application Details</b>      | Western blot, 0.1-0.5 µg/ml, Human                                                                                                                                                                               |
| <b>Subcellular Localization</b> | Membrane; Single-pass type I membrane protein. Secreted. Shedding of the ectodomain produces a soluble form.                                                                                                     |
| <b>Source</b>                   | Eukaryota                                                                                                                                                                                                        |
| <b>Contents</b>                 | Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.                                                                                                                                              |
| <b>Immunogen</b>                | E. coli-derived human Syndecan-1/SDC1 recombinant protein (Position: Q23-G254). Human Syndecan-1/SDC1 shares 69.2% and 69.8% amino acid (aa) sequence identity with mouse and rat Syndecan-1/SDC1, respectively. |
| <b>Purification</b>             | Immunogen affinity purified.                                                                                                                                                                                     |
| <b>Cross Reactivity</b>         | No cross reactivity with other proteins.                                                                                                                                                                         |
| <b>Storage</b>                  | At -20 °C for one year. After reconstitution, at 4 °C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.                               |

## Protein Information

---

|                          |                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|--------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Name</b>              | SDC1 ( <a href="#">HGNC:10658</a> )                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Synonyms</b>          | SDC                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| <b>Function</b>          | Cell surface proteoglycan that contains both heparan sulfate and chondroitin sulfate and that links the cytoskeleton to the interstitial matrix (By similarity). Regulates exosome biogenesis in concert with SDCBP and PDCD6IP (PubMed: <a href="#">22660413</a> ). Able to induce its own expression in dental mesenchymal cells and also in the neighboring dental epithelial cells via an MSX1-mediated pathway (By similarity). |
| <b>Cellular Location</b> | Membrane; Single-pass type I membrane protein. Secreted Secreted, extracellular exosome Note=Shedding of the ectodomain produces a soluble form                                                                                                                                                                                                                                                                                      |
| <b>Tissue Location</b>   | Detected in placenta (at protein level) (PubMed:32337544). Detected in fibroblasts (at protein level) (PubMed:36213313).                                                                                                                                                                                                                                                                                                             |

## Background

Syndecan 1 is a protein which in humans is encoded by the SDC1 gene. The protein encoded by this gene is a transmembrane (type I) heparan sulfate proteoglycan and is a member of the syndecan proteoglycan family. The syndecans mediate cell binding, cell signaling, and cytoskeletal organization and syndecan receptors are required for internalization of the HIV-1 tat protein. The syndecan-1 protein functions as an integral membrane protein and participates in cell proliferation, cell migration and cell-matrix interactions via its receptor for extracellular matrix proteins. Altered syndecan-1 expression has been detected in several different tumor types. While several transcript variants may exist for this gene, the full-length nature of only two have been described to date. These two represent the major variants of this gene and encode the same protein.

## Images




Figure 1. Western blot analysis of Syndecan-1/SDC1 using anti-Syndecan-1/SDC1 antibody (AB012911).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.