

Anti-NOV/CCN3 Picoband Antibody

Catalog # ABO13050

Product Information

Application	WB, IHC-P
Primary Accession	P48745
Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Format	Lyophilized
Description	Rabbit IgG polyclonal antibody for Protein NOV homolog(NOV) detection. Tested with WB, IHC-P in Human;Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Additional Information

Gene ID	4856
Other Names	Protein NOV homolog, NovH, CCN family member 3, Insulin-like growth factor-binding protein 9, IBP-9, IGF-binding protein 9, IGFBP-9, Nephroblastoma-overexpressed gene protein homolog, NOV, CCN3, IGFBP9, NOVH
Calculated MW	39162
Application Details	Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, Rat, By Heat Western blot, 0.1-0.5 μ g/ml, Human,
Subcellular Localization	Secreted.
Tissue Specificity	Expressed in bone marrow, thymic cells and nephroblastoma. Increased expression in Wilms tumor of the stromal type. .
Source	Eukaryota
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ N.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human NOV/CCN3 (334-357aa HTNCPKNNEAFLQELELKTTTRGKM), different from the related mouse sequence by seven amino acids, and from the related rat sequence by four amino acids.
Purification	Immunogen affinity purified.

Cross Reactivity	No cross reactivity with other proteins.
Storage	At -20 °C for one year. After r ° Constitution, at 4 °C for one month. It ° Can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

Protein Information

Name	CCN3 (HGNC:7885)
Synonyms	IGFBP9, NOV, NOVH
Function	Immediate-early protein playing a role in various cellular processes including proliferation, adhesion, migration, differentiation and survival (PubMed:12050162, PubMed:12695522, PubMed:15181016, PubMed:15611078, PubMed:21344378). Acts by binding to integrins or membrane receptors such as NOTCH1 (PubMed:12695522, PubMed:15611078, PubMed:21344378). Essential regulator of hematopoietic stem and progenitor cell function (PubMed:17463287). Inhibits myogenic differentiation through the activation of Notch-signaling pathway (PubMed:12050162). Inhibits vascular smooth muscle cells proliferation by increasing expression of cell-cycle regulators such as CDKN2B or CDKN1A independently of TGFB1 signaling (PubMed:20139355). Ligand of integrins ITGAV:ITGB3 and ITGA5:ITGB1, acts directly upon endothelial cells to stimulate pro-angiogenic activities and induces angiogenesis. In endothelial cells, supports cell adhesion, induces directed cell migration (chemotaxis) and promotes cell survival (PubMed:12695522). Also plays a role in cutaneous wound healing acting as integrin receptor ligand. Supports skin fibroblast adhesion through ITGA5:ITGB1 and ITGA6:ITGB1 and induces fibroblast chemotaxis through ITGAV:ITGB5. Seems to enhance bFGF-induced DNA synthesis in fibroblasts (PubMed:15611078). Involved in bone regeneration as a negative regulator (By similarity). Enhances the articular chondrocytic phenotype, whereas it repressed the one representing endochondral ossification (PubMed:21871891). Impairs pancreatic beta-cell function, inhibits beta-cell proliferation and insulin secretion (By similarity). Plays a role as negative regulator of endothelial pro-inflammatory activation reducing monocyte adhesion, its anti-inflammatory effects occur secondary to the inhibition of NF-kappaB signaling pathway (PubMed:21063504). Contributes to the control and coordination of inflammatory processes in atherosclerosis (By similarity). Attenuates inflammatory pain through regulation of IL1B- and TNF-induced MMP9, MMP2 and CCL2 expression. Inhibits MMP9 expression through ITGB1 engagement (PubMed:21871891). Brain osteoanabolic hormone (By similarity). Drives osteogenesis in osteochondral skeletal stem cells (PubMed:38987585). During lactation, maintains the maternal skeleton and viability of offspring (By similarity).
Cellular Location	Secreted {ECO:0000250 UniProtKB:Q64299}. Cytoplasm. Cell junction, gap junction. Note=Localizes at the gap junction in presence of GJA1. {ECO:0000250 UniProtKB:Q9QZQ5}
Tissue Location	Expressed in endothelial cells (at protein level) (PubMed:21063504). Expressed in bone marrow and thymic cells

Background

NOV (nephroblastoma overexpressed), also known as CCN3, is a matricellular protein that in humans is encoded by the NOV gene. The protein encoded by this gene is a small secreted cysteine-rich protein and a member of the CCN family of regulatory proteins. CNN family proteins associate with the extracellular

matrix and play an important role in cardiovascular and skeletal development, fibrosis and cancer development.

Images








Figure 5. IHC analysis of NOV/CCN3 using anti-NOV/CCN3 antibody (ABO13050).





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.