

Anti-Phospho-GluR1 (S845) GRIA1 Rabbit Monoclonal Antibody

Catalog # ABO13096

Product Information

Application	WB
Primary Accession	P42261
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Phospho-GluR1 (S845) GRIA1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	2890
Other Names	Glutamate receptor 1, GluR-1, AMPA-selective glutamate receptor 1, GluR-A, Glutamate receptor ionotropic, AMPA 1, GRIA1 (HGNC:4571)
Calculated MW	101506
Application Details	WB 1:500-1:1000
Subcellular Localization	Cell membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein. Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein. Cell junction, synapse, postsynaptic cell membrane, postsynaptic density. Cell projection, dendrite. Cell projection, dendritic spine. Interaction with CACNG2, CNIH2 and CNIH3 promotes cell surface expression.
Tissue Specificity	Detected in cerebellum (at protein level)..
Source	Eukaryota
Protein Name	Glutamate receptor 1
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: GFH-7
Immunogen	A synthesized peptide derived from human GluR1
Purification	Affinity-chromatography

Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
----------------	--

Protein Information

Name	GRIA1 (HGNC:4571)
-------------	-------------------------------------


Function	Ionotropic glutamate receptor that functions as a ligand- gated cation channel, gated by L-glutamate and glutamatergic agonists such as alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA), quisqualic acid, and kainic acid (PubMed:1311100, PubMed:20805473, PubMed:21172611, PubMed:28628100, PubMed:35675825). L- glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse upon entry of monovalent and divalent cations such as sodium and calcium. The receptor then desensitizes rapidly and enters in a transient inactive state, characterized by the presence of bound agonist (By similarity). In the presence of CACNG2 or CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of L- glutamate (PubMed:21172611). Resensitization is blocked by CNIH2 through interaction with CACNG8 in the CACNG8-containing AMPA receptors complex (PubMed:21172611). Calcium (Ca(2+)) permeability depends on subunits composition and, heteromeric channels containing edited GRIA2 subunit are calcium-impermeable. Also permeable to other divalents cations such as strontium(2+) and magnesium(2+) and monovalent cations such as potassium(1+) and lithium(1+) (By similarity).
-----------------	---

Cellular Location	Cell membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane {ECO:0000250 UniProtKB:P19490}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P19490}. Postsynaptic cell membrane; Multi-pass membrane protein. Postsynaptic density membrane {ECO:0000250 UniProtKB:P23818}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P23818}. Cell projection, dendrite {ECO:0000250 UniProtKB:P23818}. Cell projection, dendritic spine {ECO:0000250 UniProtKB:P23818}. Early endosome membrane {ECO:0000250 UniProtKB:P19490}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P19490}. Recycling endosome membrane {ECO:0000250 UniProtKB:P19490}; Multi-pass membrane protein {ECO:0000250 UniProtKB:P19490}. Presynapse {ECO:0000250 UniProtKB:P23818}. Synapse {ECO:0000250 UniProtKB:P23818} Note=Interaction with CACNG2, CNIH2 and CNIH3 promotes cell surface expression. Colocalizes with PDLIM4 in early endosomes. Displays a somatodendritic localization and is excluded from axons in neurons (By similarity). Localized to cone photoreceptor pedicles (By similarity) {ECO:0000250 UniProtKB:P19490, ECO:0000250 UniProtKB:P23818}
--------------------------	--

Tissue Location	Widely expressed in brain.
------------------------	----------------------------

Images

Western blot analysis of Phospho-GluR1 (S845) expression in (1) Human brain lysate treated with Lambda phosphatase lysate; (2) Human brain lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.