

Anti-Phospho-IRE1 (S724) ERN1 Rabbit Monoclonal Antibody

Catalog # ABO13130

Product Information

Application	WB
Primary Accession	O75460
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Phospho-IRE1 (S724) ERN1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.


Additional Information

Gene ID	2081
Other Names	Serine/threonine-protein kinase/endoribonuclease IRE1, Endoplasmic reticulum-to-nucleus signaling 1, Inositol-requiring protein 1, hIRE1p, Ire1-alpha, IRE1a, Serine/threonine-protein kinase, 2.7.11.1, Endoribonuclease, 3.1.26.-, ERN1 (HGNC:3449)
Calculated MW	109735
Application Details	WB 1:500-1:2000
Subcellular Localization	Endoplasmic reticulum membrane ; Single-pass type I membrane protein.
Tissue Specificity	Ubiquitously expressed. High levels observed in pancreatic tissue..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: EGH-5
Immunogen	A synthesized peptide derived from human Phospho-IRE1 (S724)
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Name	ERN1 (HGNC:3449)
Function	Serine/threonine-protein kinase and endoribonuclease that acts as a key sensor for the endoplasmic reticulum unfolded protein response (UPR) (PubMed: 11175748 , PubMed: 11779464 , PubMed: 12637535 , PubMed: 19328063 , PubMed: 21317875 , PubMed: 28128204 , PubMed: 30118681 , PubMed: 36739529 , PubMed: 9637683). In unstressed cells, the endoplasmic reticulum luminal domain is maintained in its inactive monomeric state by binding to the endoplasmic reticulum chaperone HSPA5/BiP (PubMed: 21317875). Accumulation of misfolded proteins in the endoplasmic reticulum causes release of HSPA5/BiP, allowing the luminal domain to homodimerize, promoting autophosphorylation of the kinase domain and subsequent activation of the endoribonuclease activity (PubMed: 21317875). The endoribonuclease activity is specific for XBP1 mRNA and excises 26 nucleotides from XBP1 mRNA (PubMed: 11779464 , PubMed: 21317875 , PubMed: 24508390). The resulting spliced transcript of XBP1 encodes a transcriptional activator protein that up-regulates expression of UPR target genes (PubMed: 11779464 , PubMed: 21317875 , PubMed: 24508390). Acts as an upstream signal for ER stress-induced GORASP2-mediated unconventional (ER/Golgi-independent) trafficking of CFTR to cell membrane by modulating the expression and localization of SEC16A (PubMed: 21884936 , PubMed: 28067262).
Cellular Location	Endoplasmic reticulum membrane; Single-pass type I membrane protein
Tissue Location	Ubiquitously expressed. High levels observed in pancreatic tissue.

Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.

Western blot analysis of Phospho-IRE1 (S724) expression in (1) K562 cell lysate treated with AP; (2) Untreated K562 cell lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.