

Anti-Phospho-c-Myc (T58) Rabbit Monoclonal Antibody

Catalog # ABO13131

Product Information

Application	WB, IF, ICC, FC
Primary Accession	P01106
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Phospho-c-Myc (T58) Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.

Additional Information

Gene ID	4609
Other Names	Myc proto-oncogene protein, Class E basic helix-loop-helix protein 39, bHLHe39, Proto-oncogene c-Myc, Transcription factor p64, MYC, BHLHE39
Calculated MW	50565
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 FC 1:50
Subcellular Localization	Nucleus, nucleoplasm. Nucleus, nucleolus.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: FOC-13
Immunogen	A synthesized peptide derived from human Phospho-c-Myc (T58)
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	MYC
Synonyms	BHLHE39


Function

Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3' (PubMed:[24940000](#), PubMed:[25956029](#)). Activates the transcription of growth-related genes (PubMed:[24940000](#), PubMed:[25956029](#)). Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed:[24940000](#), PubMed:[25956029](#)). Regulator of somatic reprogramming, controls self-renewal of embryonic stem cells (By similarity). Functions with TAF6L to activate target gene expression through RNA polymerase II pause release (By similarity). Positively regulates transcription of HNRNPA1, HNRNPA2 and PTBP1 which in turn regulate splicing of pyruvate kinase PKM by binding repressively to sequences flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production of the PKM M2 isoform (PubMed:[20010808](#)).

Cellular Location

Nucleus, nucleoplasm. Nucleus, nucleolus. Nucleus. Cytoplasm Chromosome. Note=Association with chromatin is reduced by hyperphosphorylation (PubMed:30158517) Localization to the nucleolus is dependent on HEATR1 (PubMed:38225354)

Images



Western blot analysis of Phospho-c-Myc (T58) expression in (1) HeLa cell lysate; (2) HeLa cell lysate treated with Calyculin A and Okadaic Acid.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.