

Anti-Phospho-PLK1 (T210) Rabbit Monoclonal Antibody

Catalog # ABO13138

Product Information

Application	WB
Primary Accession	P53350
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Phospho-PLK1 (T210) Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.

Additional Information

Gene ID	5347
Other Names	Serine/threonine-protein kinase PLK1, 2.7.11.21, Polo-like kinase 1, PLK-1, Serine/threonine-protein kinase 13, STPK13, PLK1, PLK
Calculated MW	68255
Application Details	WB 1:1000-1:2000
Subcellular Localization	Nucleus. Chromosome, centromere, kinetochore. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Midbody. During early stages of mitosis, the phosphorylated form is detected on centrosomes and kinetochores. Localizes to the outer kinetochore. Presence of SGOL1 and interaction with the phosphorylated form of BUB1 is required for the kinetochore localization. Localizes onto the central spindle by phosphorylating and docking at midzone proteins KIF20A/MKLP2 and PRC1. Colocalizes with FRY to separating centrosomes and spindle poles from prophase to metaphase in mitosis, but not in other stages of the cell cycle.
Tissue Specificity	Placenta and colon.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: HAO-16
Immunogen	A synthesized peptide derived from human PLK1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at

4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name PLK1

Synonyms PLK

Function Serine/threonine-protein kinase that performs several important functions throughout M phase of the cell cycle, including the regulation of centrosome maturation and spindle assembly, the removal of cohesins from chromosome arms, the inactivation of anaphase-promoting complex/cyclosome (APC/C) inhibitors, and the regulation of mitotic exit and cytokinesis (PubMed:[11202906](#), PubMed:[12207013](#), PubMed:[12447691](#), PubMed:[12524548](#), PubMed:[12738781](#), PubMed:[12852856](#), PubMed:[12939256](#), PubMed:[14532005](#), PubMed:[14734534](#), PubMed:[15070733](#), PubMed:[15148369](#), PubMed:[15469984](#), PubMed:[16198290](#), PubMed:[16247472](#), PubMed:[16980960](#), PubMed:[17081991](#), PubMed:[17351640](#), PubMed:[17376779](#), PubMed:[17617734](#), PubMed:[18174154](#), PubMed:[18331714](#), PubMed:[18418051](#), PubMed:[18477460](#), PubMed:[18521620](#), PubMed:[18615013](#), PubMed:[19160488](#), PubMed:[19351716](#), PubMed:[19468300](#), PubMed:[19468302](#), PubMed:[19473992](#), PubMed:[19509060](#), PubMed:[19597481](#), PubMed:[23455478](#), PubMed:[23509069](#), PubMed:[28512243](#), PubMed:[8991084](#)). Polo-like kinase proteins act by binding and phosphorylating proteins that are already phosphorylated on a specific motif recognized by the POLO box domains (PubMed:[11202906](#), PubMed:[12207013](#), PubMed:[12447691](#), PubMed:[12524548](#), PubMed:[12738781](#), PubMed:[12852856](#), PubMed:[12939256](#), PubMed:[14532005](#), PubMed:[14734534](#), PubMed:[15070733](#), PubMed:[15148369](#), PubMed:[15469984](#), PubMed:[16198290](#), PubMed:[16247472](#), PubMed:[16980960](#), PubMed:[17081991](#), PubMed:[17351640](#), PubMed:[17376779](#), PubMed:[17617734](#), PubMed:[18174154](#), PubMed:[18331714](#), PubMed:[18418051](#), PubMed:[18477460](#), PubMed:[18521620](#), PubMed:[18615013](#), PubMed:[19160488](#), PubMed:[19351716](#), PubMed:[19468300](#), PubMed:[19468302](#), PubMed:[19473992](#), PubMed:[19509060](#), PubMed:[19597481](#), PubMed:[23455478](#), PubMed:[23509069](#), PubMed:[28512243](#), PubMed:[8991084](#)). Phosphorylates BORA, BUB1B/BUBR1, CCNB1, CDC25C, CEP55, ECT2, ERCC6L, FBXO5/EMI1, FOXM1, KIF20A/MKLP2, CENPU, NEDD1, NINL, NPM1, NUDC, PKMYT1/MYT1, KIZ, MRE11, PPP1R12A/MYPT1, POLQ, PRC1, RACGAP1/CYK4, RAD51, RHNO1, SGO1, STAG2/SA2, TEX14, TOPORS, p73/TP73, TPT1, WEE1 and HNRNPU (PubMed:[11202906](#), PubMed:[12207013](#), PubMed:[12447691](#), PubMed:[12524548](#), PubMed:[12738781](#), PubMed:[12852856](#), PubMed:[12939256](#), PubMed:[14532005](#), PubMed:[14734534](#), PubMed:[15070733](#), PubMed:[15148369](#), PubMed:[15469984](#), PubMed:[16198290](#), PubMed:[16247472](#), PubMed:[16980960](#), PubMed:[17081991](#), PubMed:[17218258](#), PubMed:[17351640](#), PubMed:[17376779](#), PubMed:[17617734](#), PubMed:[18174154](#), PubMed:[18331714](#), PubMed:[18418051](#), PubMed:[18477460](#), PubMed:[18521620](#), PubMed:[18615013](#), PubMed:[19160488](#), PubMed:[19351716](#), PubMed:[19468300](#), PubMed:[19468302](#), PubMed:[19473992](#), PubMed:[19509060](#), PubMed:[19597481](#), PubMed:[22325354](#), PubMed:[23455478](#), PubMed:[23509069](#), PubMed:[25986610](#), PubMed:[26811421](#), PubMed:[28512243](#), PubMed:[37440612](#), PubMed:[37674080](#), PubMed:[8991084](#)). Plays a key role in centrosome functions and the assembly of bipolar spindles by phosphorylating KIZ, NEDD1 and NINL (PubMed:[16980960](#),

PubMed:[19509060](#)). NEDD1 phosphorylation promotes subsequent targeting of the gamma-tubulin ring complex (gTuRC) to the centrosome, an important step for spindle formation (PubMed:[19509060](#)). Phosphorylation of NINL component of the centrosome leads to NINL dissociation from other centrosomal proteins (PubMed:[12852856](#)). Involved in mitosis exit and cytokinesis by phosphorylating CEP55, ECT2, KIF20A/MKLP2, CENPU, PRC1 and RACGAP1 (PubMed:[12939256](#), PubMed:[16247472](#), PubMed:[17351640](#), PubMed:[19468300](#), PubMed:[19468302](#)). Recruited at the central spindle by phosphorylating and docking PRC1 and KIF20A/MKLP2; creates its own docking sites on PRC1 and KIF20A/MKLP2 by mediating phosphorylation of sites subsequently recognized by the POLO box domains (PubMed:[12939256](#), PubMed:[17351640](#)). Phosphorylates RACGAP1, thereby creating a docking site for the Rho GTP exchange factor ECT2 that is essential for the cleavage furrow formation (PubMed:[19468300](#), PubMed:[19468302](#)). Promotes the central spindle recruitment of ECT2 (PubMed:[16247472](#)). Plays a central role in G2/M transition of mitotic cell cycle by phosphorylating CCNB1, CDC25C, FOXM1, CENPU, PKMYT1/MYT1, PPP1R12A/MYPT1 and WEE1 (PubMed:[11202906](#), PubMed:[12447691](#), PubMed:[12524548](#), PubMed:[19160488](#)). Part of a regulatory circuit that promotes the activation of CDK1 by phosphorylating the positive regulator CDC25C and inhibiting the negative regulators WEE1 and PKMYT1/MYT1 (PubMed:[11202906](#)). Also acts by mediating phosphorylation of cyclin-B1 (CCNB1) on centrosomes in prophase (PubMed:[12447691](#), PubMed:[12524548](#)). Phosphorylates FOXM1, a key mitotic transcription regulator, leading to enhance FOXM1 transcriptional activity (PubMed:[19160488](#)). Involved in kinetochore functions and sister chromatid cohesion by phosphorylating BUB1B/BUBR1, FBXO5/EMI1 and STAG2/SA2 (PubMed:[15148369](#), PubMed:[15469984](#), PubMed:[17376779](#), PubMed:[18331714](#)). PLK1 is high on non-attached kinetochores suggesting a role of PLK1 in kinetochore attachment or in spindle assembly checkpoint (SAC) regulation (PubMed:[17617734](#)). Required for kinetochore localization of BUB1B (PubMed:[17376779](#)). Regulates the dissociation of cohesin from chromosomes by phosphorylating cohesin subunits such as STAG2/SA2 (By similarity). Phosphorylates SGO1: required for spindle pole localization of isoform 3 of SGO1 and plays a role in regulating its centriole cohesion function (PubMed:[18331714](#)). Mediates phosphorylation of FBXO5/EMI1, a negative regulator of the APC/C complex during prophase, leading to FBXO5/EMI1 ubiquitination and degradation by the proteasome (PubMed:[15148369](#), PubMed:[15469984](#)). Acts as a negative regulator of p53 family members: phosphorylates TOPORS, leading to inhibit the sumoylation of p53/TP53 and simultaneously enhance the ubiquitination and subsequent degradation of p53/TP53 (PubMed:[19473992](#)). Phosphorylates the transactivation domain of the transcription factor p73/TP73, leading to inhibit p73/TP73-mediated transcriptional activation and pro-apoptotic functions. Phosphorylates BORA, and thereby promotes the degradation of BORA (PubMed:[18521620](#)). Contributes to the regulation of AURKA function (PubMed:[18615013](#), PubMed:[18662541](#)). Also required for recovery after DNA damage checkpoint and entry into mitosis (PubMed:[18615013](#), PubMed:[18662541](#)). Phosphorylates MISP, leading to stabilization of cortical and astral microtubule attachments required for proper spindle positioning (PubMed:[23509069](#)). Together with MEIKIN, acts as a regulator of kinetochore function during meiosis I: required both for mono- orientation of kinetochores on sister chromosomes and protection of centromeric cohesin from separase-mediated cleavage (By similarity). Phosphorylates CEP68 and is required for its degradation (PubMed:[25503564](#)). Regulates nuclear envelope breakdown during prophase by phosphorylating DCTN1 resulting in its localization in the nuclear envelope (PubMed:[20679239](#)). Phosphorylates the heat shock transcription factor HSF1, promoting HSF1 nuclear translocation upon heat shock (PubMed:[15661742](#)). Phosphorylates HSF1 also in the early mitotic period; this phosphorylation regulates HSF1 localization to the spindle pole, the recruitment of the SCF(BTRC) ubiquitin ligase complex inducing

HSF1 degradation, and hence mitotic progression (PubMed:[18794143](#)). Regulates mitotic progression by phosphorylating RIOK2 (PubMed:[21880710](#)). Through the phosphorylation of DZIP1 regulates the localization during mitosis of the BBSome, a ciliary protein complex involved in cilium biogenesis (PubMed:[27979967](#)). Regulates DNA repair during mitosis by mediating phosphorylation of POLQ and RHNO1, thereby promoting POLQ recruitment to DNA damage sites (PubMed:[37440612](#), PubMed:[37674080](#)). Phosphorylates ATXN10 which may play a role in the regulation of cytokinesis and may stimulate the proteasome-mediated degradation of ATXN10 (PubMed:[21857149](#)).


Cellular Location

Nucleus. Chromosome, centromere, kinetochore. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Midbody. Note=localization at the centrosome starts at the G1/S transition (PubMed:24018379). During early stages of mitosis, the phosphorylated form is detected on centrosomes and kinetochores. Localizes to the outer kinetochore. Presence of SGO1 and interaction with the phosphorylated form of BUB1 is required for the kinetochore localization. Localizes onto the central spindle by phosphorylating and docking at midzone proteins KIF20A/MKLP2 and PRC1 Colocalizes with FRY to separating centrosomes and spindle poles from prophase to metaphase in mitosis, but not in other stages of the cell cycle. Localization to the centrosome is required for S phase progression (PubMed:24018379). Colocalizes with HSF1 at the spindle poles during prometaphase (PubMed:18794143)

Tissue Location

Placenta and colon.

Images



Western blot analysis of Phospho-PLK1 (T210) expression in (1) HeLa cell lysate; (2) HeLa cell treated with nocodazole lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.