

Anti-Phospho-SHIP (Y1020) INPP5D Rabbit Monoclonal Antibody

Catalog # ABO13170

Product Information

Application	WB, IF, ICC, IP
Primary Accession	Q92835
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Phospho-SHIP (Y1020) INPP5D Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP applications. This antibody reacts with Human.

Additional Information


Gene ID	3635
Other Names	Phosphatidylinositol 3, 4, 5-trisphosphate 5-phosphatase 1, 3.1.3.86, Inositol polyphosphate-5-phosphatase D, 3.1.3.56, Inositol polyphosphate-5-phosphatase of 145 kDa, SIP-145, Phosphatidylinositol 4, 5-bisphosphate 5-phosphatase, 3.1.3.36, SH2 domain-containing inositol 5'-phosphatase 1, SH2 domain-containing inositol phosphatase 1, SHIP-1, p150Ship, hp51CN, INPP5D, SHIP {ECO:0000303 PubMed:10764818}, SHIP1
Calculated MW	133292
Application Details	WB 1:1000-1:2000 ICC/IF 1:50-1:200 IP 1:20
Subcellular Localization	Cytoplasm. Cell membrane ; Peripheral membrane protein. Membrane raft. Cytoplasm, cytoskeleton. Membrane ; Peripheral membrane protein. Translocates to the plasma membrane when activated, translocation is probably due to different mechanisms depending on the stimulus and cell type. Translocates from the cytoplasm to membrane ruffles in a FCGR3/CD16-dependent manner. Colocalizes with FC-gamma-RIIB receptor (FCGR2B) or FCGR3/CD16 at membrane ruffles. Tyrosine phosphorylation may also participate in membrane localization..
Tissue Specificity	Specifically expressed in immune and hematopoietic cells. Expressed in bone marrow and blood cells. Levels vary considerably within this compartment. Present in at least 74% of immature CD34+ cells, whereas within the more mature population of CD33+ cells, it is present in only 10% of cells. Present in the majority of T-cells, while it is present in a minority of B-cells (at protein level)..

Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: HFF-9 A synthesized peptide derived from human SHIP
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	INPP5D
Synonyms	SHIP {ECO:0000303 PubMed:10764818}, SHIP
Function	Phosphatidylinositol (PtdIns) phosphatase that specifically hydrolyzes the 5-phosphate of phosphatidylinositol-3,4,5-trisphosphate (PtdIns(3,4,5)P3) to produce PtdIns(3,4)P2, thereby negatively regulating the PI3K (phosphoinositide 3-kinase) pathways (PubMed:10764818, PubMed:8723348, PubMed:8769125). Also, is able to hydrolyze the 5-phosphate of phosphatidylinositol-4,5-bisphosphate (PtdIns(4,5)P3), 1D-myo-inositol 1,4,5-trisphosphate and inositol 1,3,4,5-tetrakisphosphate (PubMed:10764818, PubMed:40969890, PubMed:8769125, PubMed:9108392). Acts as a negative regulator of B-cell antigen receptor signaling. Mediates signaling from the FC-gamma-RIIB receptor (FCGR2B), playing a central role in terminating signal transduction from activating immune/hematopoietic cell receptor systems. Acts as a negative regulator of myeloid cell proliferation/survival and chemotaxis, mast cell degranulation, immune cells homeostasis, integrin alpha-IIb/beta-3 signaling in platelets and JNK signaling in B-cells. Regulates proliferation of osteoclast precursors, macrophage programming, phagocytosis and activation and is required for endotoxin tolerance. Involved in the control of cell-cell junctions, CD32a signaling in neutrophils and modulation of EGF-induced phospholipase C activity (PubMed:16682172). Key regulator of neutrophil migration, by governing the formation of the leading edge and polarization required for chemotaxis. Modulates FCGR3/CD16-mediated cytotoxicity in NK cells. Mediates the activin/TGF-beta-induced apoptosis through its Smad-dependent expression.
Cellular Location	Cytoplasm. Cell membrane {ECO:0000250 UniProtKB:Q9ES52}; Peripheral membrane protein {ECO:0000250 UniProtKB:Q9ES52}. Membrane raft {ECO:0000250 UniProtKB:Q9ES52}. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:Q9ES52}. Membrane; Peripheral membrane protein Note=Translocates to the plasma membrane when activated, translocation is probably due to different mechanisms depending on the stimulus and cell type. Translocates from the cytoplasm to membrane ruffles in a FCGR3/CD16-dependent manner. Colocalizes with FC-gamma-RIIB receptor (FCGR2B) or FCGR3/CD16 at membrane ruffles. Tyrosine phosphorylation may also participate in membrane localization {ECO:0000250 UniProtKB:Q9ES52}
Tissue Location	Specifically expressed in immune and hematopoietic cells. Expressed in bone marrow and blood cells. Levels vary considerably within this compartment. Present in at least 74% of immature CD34+ cells, whereas within the more mature population of CD33+ cells, it is present in only 10% of cells. Present in the majority of T-cells, while it is present in a minority of B-cells (at protein level).

Images



Western blot analysis of Phospho-SHIP (Y1020) expression in (1) Raji cell lysate; (2) Raji cell treated with pervanadate lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.