

Anti-C9/Complement Component C9 Rabbit Monoclonal Antibody

Catalog # ABO13214

Product Information

Application	WB, IHC, IP
Primary Accession	P02748
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-C9/Complement Component C9 Rabbit Monoclonal Antibody . Tested in WB, IHC, IP applications. This antibody reacts with Human.


Additional Information

Gene ID	735
Other Names	Complement component C9, Complement component C9a, Complement component C9b, C9
Calculated MW	63173
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 IP 1:50
Subcellular Localization	Secreted. Cell membrane; Multi-pass membrane protein. Secreted as soluble monomer. Oligomerizes at target membranes, forming a pre-pore. A conformation change then leads to the formation of a 100 Angstrom diameter pore.
Tissue Specificity	Plasma.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: BAO-3
Immunogen	A synthesized peptide derived from human C9
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	C9 {ECO:0000303 PubMed:4018030, ECO:0000312 HGNC:HGNC:1358}
Function	Pore-forming component of the membrane attack complex (MAC), a multiprotein complex activated by the complement cascade, which inserts into a target cell membrane and forms a pore, leading to target cell membrane rupture and cell lysis (PubMed:22832194, PubMed:26841837, PubMed:26841934, PubMed:27052168, PubMed:30552328, PubMed:6177822, PubMed:9212048, PubMed:9634479). The MAC is initiated by proteolytic cleavage of C5 into complement C5b in response to the classical, alternative, lectin and GZMK complement pathways (PubMed:39914456, PubMed:39814882, PubMed:9212048, PubMed:9634479). The complement pathways consist in a cascade of proteins that leads to phagocytosis and breakdown of pathogens and signaling that strengthens the adaptive immune system (PubMed:9212048, PubMed:9634479). Constitutes the pore-forming subunit of the MAC complex: during MAC assembly, C9 associates with the C5b8 intermediate complex, and polymerizes to complete the pore (PubMed:26841934, PubMed:30111885, PubMed:30552328, PubMed:34752492, PubMed:4055801, PubMed:6177822).
Cellular Location	Secreted. Target cell membrane; Multi-pass membrane protein. Note=Secreted as soluble monomer (PubMed:26841934, PubMed:30111885, PubMed:4055801, PubMed:9634479) Oligomerizes at target membranes, forming a pre-pore (PubMed:26841934, PubMed:30111885, PubMed:31061395, PubMed:4055801, PubMed:9634479). A conformation change then leads to the formation of a 100 Angstrom diameter pore (PubMed:26841934, PubMed:30111885, PubMed:31061395, PubMed:4055801, PubMed:9634479).
Tissue Location	Plasma (at protein level).

Images



Western blot analysis of C9 expression in Human fetal liver lysate.

Immunohistochemical analysis of paraffin-embedded human kidney, using C9 Antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.