

Anti-ILK/Integrin Linked Ilk Rabbit Monoclonal Antibody

Catalog # ABO13259

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	Q13418
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-ILK/Integrin Linked Ilk Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	3611
Other Names	Integrin-linked protein kinase, 2.7.11.1, 59 kDa serine/threonine-protein kinase, Beta-integrin-linked kinase, ILK-1, ILK-2, p59ILK, ILK (HGNC:6040)
Calculated MW	51419
Application Details	WB 1:1000-1:5000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Subcellular Localization	Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium. Cytoplasm, myofibril, sarcomere.
Tissue Specificity	Highly expressed in heart followed by skeletal muscle, pancreas and kidney. Weakly expressed in placenta, lung and liver.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: FHE-9
Immunogen	A synthesized peptide derived from human ILK
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at

4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information



Name	ILK (HGNC:6040)
Function	Scaffold protein which mediates protein-protein interactions during a range of cellular events including focal adhesion assembly, cell adhesion and cell migration (PubMed:17420447, PubMed:20005845, PubMed:30367047, PubMed:32528174). Regulates integrin-mediated signal transduction by contributing to inside-out integrin activation (By similarity). Recruits PARVA and LIMS1/PITCH to form the heterotrimeric IPP (ILK-PINCH-PARVIN) complex which binds to F-actin via the C- terminal tail of LIMS1 and the N-terminal region of PARVA, promoting F- actin filament bundling, a process required to generate force for actin cytoskeleton reorganization and subsequent dynamic cell adhesion events such as cell spreading and migration (PubMed:30367047). Binding to PARVA promotes effective assembly of ILK into focal adhesions while PARVA-bound ILK can simultaneously engage integrin-beta cytoplasmic tails to mediate cell adhesion (PubMed:20005845). Plays a role with PARVG in promoting the cell adhesion and spreading of leukocytes (PubMed:16517730). Acts as an upstream effector of both AKT1/PKB and GSK3 (PubMed:9736715). Mediates trafficking of caveolae to the cell surface in an ITGB1-dependent manner by promoting the recruitment of IQGAP1 to the cell cortex which cooperates with its effector DIAPH1 to locally stabilize microtubules and allow stable insertion of caveolae into the plasma membrane (By similarity). Required for the maintenance of mitotic spindle integrity by promoting phosphorylation of TACC3 by AURKA (PubMed:18283114). Associates with chromatin and may act as a negative regulator of transcription when located in the nucleus (PubMed:17420447).
Cellular Location	Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium {ECO:0000250 UniProtKB:O55222}. Cytoplasm, myofibril, sarcomere. Cytoplasm Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cell cortex {ECO:0000250 UniProtKB:O55222}
Tissue Location	Highly expressed in heart followed by skeletal muscle, pancreas and kidney. Weakly expressed in placenta, lung and liver

Images



Western blot analysis of ILK expression in K562 cell lysate.

Immunohistochemical analysis of paraffin-embedded human stomach, using ILK Antibody.



Immunofluorescent analysis of 293 cells, using ILK Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.