

Anti-MIF Rabbit Monoclonal Antibody

Catalog # ABO13288

Product Information

Application	WB, IP, FC
Primary Accession	P14174
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-MIF Rabbit Monoclonal Antibody . Tested in WB, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Gene ID	4282
Other Names	Macrophage migration inhibitory factor, MIF, 5.3.2.1, Glycosylation-inhibiting factor, L-dopachrome isomerase, L-dopachrome tautomerase, 5.3.3.12, Phenylpyruvate tautomerase, MIF {ECO:0000303 PubMed:2552447, ECO:0000312 HGNC:HGNC:7097}
Calculated MW	12476
Application Details	WB 1:500-1:2000 IP 1:50 FC 1:50
Subcellular Localization	Secreted. Cytoplasm. Does not have a cleavable signal sequence and is secreted via a specialized, non- classical pathway. Secreted by macrophages upon stimulation by bacterial lipopolysaccharide (LPS), or by M.tuberculosis antigens.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ABAG-13
Immunogen	A synthesized peptide derived from human MIF
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	MIF {ECO:0000303 PubMed:2552447, ECO:0000312 HGNC:HGNC:7097}
Function	Pro-inflammatory cytokine involved in the innate immune response to bacterial pathogens (PubMed: 15908412 , PubMed: 17443469 , PubMed: 23776208). The expression of MIF at sites of inflammation suggests a role as mediator in regulating the function of macrophages in host defense (PubMed: 15908412 , PubMed: 17443469 , PubMed: 23776208). Counteracts the anti-inflammatory activity of glucocorticoids (PubMed: 15908412 , PubMed: 17443469 , PubMed: 23776208). Has phenylpyruvate tautomerase and dopachrome tautomerase activity (in vitro), but the physiological substrate is not known (PubMed: 11439086 , PubMed: 17526494). It is not clear whether the tautomerase activity has any physiological relevance, and whether it is important for cytokine activity (PubMed: 11439086 , PubMed: 17526494).
Cellular Location	Secreted. Cytoplasm. Note=Does not have a cleavable signal sequence and is secreted via a specialized, non-classical pathway Secreted by macrophages upon stimulation by bacterial lipopolysaccharide (LPS), or by M.tuberculosis antigens

Images



Western blot analysis of MIF expression in (1) THP-1 cell lysate; (2) Mouse brain lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.