

Anti-LOX Rabbit Monoclonal Antibody

Catalog # ABO13289

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P28300
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-LOX Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	4015
Other Names	Protein-lysine 6-oxidase, 1.4.3.13, Lysyl oxidase, Protein-lysine 6-oxidase, long form, Protein-lysine 6-oxidase, short form, LOX
Calculated MW	46944
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:60
Subcellular Localization	Secreted, extracellular space.
Tissue Specificity	Heart, placenta, skeletal muscle, kidney, lung and pancreas..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ABEO-12
Immunogen	A synthesized peptide derived from human LOX
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	LOX
Function	Responsible for the post-translational oxidative deamination of peptidyl lysine residues in precursors to fibrous collagen and elastin (PubMed: 26838787). Regulator of Ras expression. May play a role in tumor suppression. Plays a role in the aortic wall architecture (By similarity).
Cellular Location	Secreted. Secreted, extracellular space
Tissue Location	Heart, placenta, skeletal muscle, kidney, lung and pancreas.

Images




Figure 1. Western blot analysis of LOX using anti-LOX antibody (M00575).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human SiHa whole cell lysates,

Lane 2: rat brain tissue lysates,

Lane 3: mouse brain tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-LOX antigen affinity purified monoclonal antibody (Catalog # M00575) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for LOX at approximately 47 kDa. The expected band size for LOX is at 47 kDa.





Figure 2. IHC analysis of LOX using anti-LOX antibody (M00575).

LOX was detected in a paraffin-embedded section of human colorectal adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-LOX Antibody (M00575) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Figure 3. IHC analysis of LOX using anti-LOX antibody (M00575).

LOX was detected in a paraffin-embedded section of human lung squamous cell carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was



blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-LOX Antibody (M00575) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.





Figure 4. IHC analysis of LOX using anti-LOX antibody (M00575).

LOX was detected in a paraffin-embedded section of human spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-LOX Antibody (M00575) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



Immunofluorescent analysis of Jurkat cells, using LOX Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.