

Anti-BMP4 Rabbit Monoclonal Antibody

Catalog # ABO13325

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P12644
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-BMP4 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.


Additional Information

Gene ID	652
Other Names	Bone morphogenetic protein 4 {ECO:0000312 HGNC:HGNC:1071}, BMP-4, BMP4 (HGNC:1071)
Calculated MW	46555
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Subcellular Localization	Secreted, extracellular space, extracellular matrix.
Tissue Specificity	Expressed in the lung and lower levels seen in the kidney. Present also in normal and neoplastic prostate tissues, and prostate cancer cell lines.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: BBB-2
Immunogen	A synthesized peptide derived from human BMP4
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	BMP4 (HGNC:1071)
Function	Growth factor of the TGF-beta superfamily that plays essential roles in many developmental processes, including neurogenesis, vascular development, angiogenesis and osteogenesis (PubMed: 31363885). Acts in concert with PTHLH/PTHRP to stimulate ductal outgrowth during embryonic mammary development and to inhibit hair follicle induction (By similarity). Initiates the canonical BMP signaling cascade by associating with type I receptor BMPR1A and type II receptor BMPR2 (PubMed: 25868050 , PubMed: 8006002). Once all three components are bound together in a complex at the cell surface, BMPR2 phosphorylates and activates BMPR1A. In turn, BMPR1A propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target genes (PubMed: 25868050 , PubMed: 29212066). Positively regulates the expression of odontogenic development regulator MSX1 via inducing the IPO7- mediated import of SMAD1 to the nucleus (By similarity). Required for MSX1-mediated mesenchymal molar tooth bud development beyond the bud stage, via promoting Wnt signaling (By similarity). Acts as a positive regulator of odontoblast differentiation during mesenchymal tooth germ formation, expression is repressed during the bell stage by MSX1- mediated inhibition of CTNNB1 signaling (By similarity). Able to induce its own expression in dental mesenchymal cells and also in the neighboring dental epithelial cells via an MSX1-mediated pathway (By similarity). Can also signal through non-canonical BMP pathways such as ERK/MAP kinase, PI3K/Akt, or SRC cascades (PubMed: 31363885). For example, induces SRC phosphorylation which, in turn, activates VEGFR2, leading to an angiogenic response (PubMed: 31363885).
Cellular Location	Secreted, extracellular space, extracellular matrix
Tissue Location	Expressed in the lung and lower levels seen in the kidney. Present also in normal and neoplastic prostate tissues, and prostate cancer cell lines

Images



Immunohistochemical analysis of paraffin-embedded human breast cancer, using BMP4 Antibody(M00321) BMP4 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-BMP4 Antibody (M00321)overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

Figure 1. Western blot analysis of BMP4 using anti-BMP4 antibody (M00321). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of



sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human COLO320 whole cell lysates,

Lane 3: human CACO-2 whole cell lysates.

After electrophoresis, proteins were transferred to a

nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5

hour at RT. The membrane was incubated with rabbit

anti-BMP4 antigen affinity purified monoclonal antibody

(Catalog # M00321) at 1:1000 overnight at 4°C, then

washed with TBS-0.1%Tween 3 times with 5 minutes each

and probed with a goat anti-rabbit IgG-HRP secondary

antibody at a dilution of 1:1000 for 1.5 hour at RT. The

signal is developed using an Enhanced Chemiluminescent

detection (ECL) kit (Catalog # EK1002) with Tanon 5200

system. A specific band was detected for BMP4 at

approximately 47 kDa. The expected band size for BMP4

is at 47 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.