

Anti-LRP1/Lrp 1 Cluster Ii Rabbit Monoclonal Antibody

Catalog # ABO13332

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	Q07954
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-LRP1/Lrp 1 Cluster Ii Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	4035
Other Names	Prolow-density lipoprotein receptor-related protein 1, LRP-1, Alpha-2-macroglobulin receptor, A2MR, Apolipoprotein E receptor, APOER, CD91, Low-density lipoprotein receptor-related protein 1 85 kDa subunit, LRP-85, Low-density lipoprotein receptor-related protein 1 515 kDa subunit, LRP-515, Low-density lipoprotein receptor-related protein 1 intracellular domain, LRPICD, LRP1 (HGNC:6692), A2MR, APR
Calculated MW	504606
Application Details	WB 1:10000-1:20000 IHC 1:50-1:100 ICC/IF 1:50-1:100 IP 1:30 FC 1:50
Subcellular Localization	Low-density lipoprotein receptor-related protein 1 85 kDa subunit: Cell membrane; Single-pass type I membrane protein. Membrane, coated pit.
Tissue Specificity	Most abundant in liver, brain and lung.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: BDC-12
Immunogen	A synthesized peptide derived from human LRP1
Purification	Affinity-chromatography

Storage

Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name

LRP1 ([HGNC:6692](#))

Synonyms

A2MR, APR

Function

Endocytic receptor involved in endocytosis and in phagocytosis of apoptotic cells (PubMed:[11907044](#), PubMed:[12713657](#)). Required for early embryonic development (By similarity). Involved in cellular lipid homeostasis. Involved in the plasma clearance of chylomicron remnants and activated LRPAP1 (alpha 2-macroglobulin), as well as the local metabolism of complexes between plasminogen activators and their endogenous inhibitors. Acts as an LRPAP1 alpha-2- macroglobulin receptor (PubMed:[1702392](#), PubMed:[26142438](#)). Acts as TAU/MAPT receptor and controls the endocytosis of TAU/MAPT as well as its subsequent spread (PubMed:[32296178](#)). May modulate cellular events, such as APP metabolism, kinase-dependent intracellular signaling, neuronal calcium signaling as well as neurotransmission (PubMed:[12888553](#)). Also acts as a receptor for IGFBP3 to mediate cell growth inhibition (PubMed:[9252371](#)).


Cellular Location

[Low-density lipoprotein receptor-related protein 1 85 kDa subunit]: Cell membrane; Single-pass type I membrane protein Membrane, coated pit [Low-density lipoprotein receptor-related protein 1 intracellular domain]: Cytoplasm Nucleus. Note=After cleavage, the intracellular domain (LRPICD) is detected both in the cytoplasm and in the nucleus.


Tissue Location

Most abundant in liver, brain and lung.


Images




Western blot analysis of LRP1 expression in A549 cell lysate.




All lanes use the Antibody at 1:3W dilution for 1 hour at room temperature.




Immunohistochemical analysis of paraffin-embedded human liver carcinoma, using LRP1 Antibody.



Immunohistochemical analysis of paraffin-embedded Human pituitary tumor, using the Antibody at 1:300 dilution.




Immunohistochemical analysis of paraffin-embedded Human prostate cancer, using the Antibody at 1:300 dilution.



Immunohistochemical analysis of paraffin-embedded Human breast cancer, using the Antibody at 1:300 dilution.

Immunofluorescent analysis using the Antibody at 1:50 dilution.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.