

Anti-LEF1 Rabbit Monoclonal Antibody

Catalog # ABO13361

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	Q9UJU2
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-LEF1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	51176
Other Names	Lymphoid enhancer-binding factor 1 {ECO:0000312 HGNC:HGNC:6551}, LEF-1, T cell-specific transcription factor 1-alpha, TCF1-alpha {ECO:0000312 HGNC:HGNC:6551}, LEF1 (HGNC:6551)
Calculated MW	44201
Application Details	WB 1:1000-1:2000 IHC 1:50-1:200 ICC/IF 1:100-1:500 IP 1:50 FC 1:50
Subcellular Localization	Nucleus. Found in nuclear bodies upon PIASG binding..
Tissue Specificity	Detected in thymus. Not detected in normal colon, but highly expressed in colon cancer biopsies and colon cancer cell lines. Expressed in several pancreatic tumors and weakly expressed in normal pancreatic tissue. Isoforms 1 and 5 are detected in several pancreatic cell lines..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: CFG-12
Immunogen	A synthesized peptide derived from human LEF1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at

4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Name	LEF1 (HGNC:6551)
Function	Transcription factor that binds DNA in a sequence-specific manner (PubMed: 2010090). Participates in the Wnt signaling pathway (By similarity). Activates transcription of target genes in the presence of CTNNB1 and EP300 (By similarity). PIAG antagonizes both Wnt-dependent and Wnt-independent activation by LEF1 (By similarity). TLE1, TLE2, TLE3 and TLE4 repress transactivation mediated by LEF1 and CTNNB1 (PubMed: 11266540). Regulates T-cell receptor alpha enhancer function (PubMed: 19653274). Required for IL17A expressing gamma-delta T-cell maturation and development, via binding to regulator loci of BLK to modulate expression (By similarity). Acts as a positive regulator of odontoblast differentiation during mesenchymal tooth germ formation, expression is repressed during the bell stage by MSX1-mediated inhibition of CTNNB1 signaling (By similarity). May play a role in hair cell differentiation and follicle morphogenesis (By similarity).
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00267}. Note=Found in nuclear bodies upon PIASG binding.
Tissue Location	Detected in thymus. Not detected in normal colon, but highly expressed in colon cancer biopsies and colon cancer cell lines. Expressed in several pancreatic tumors and weakly expressed in normal pancreatic tissue. Isoforms 1 and 5 are detected in several pancreatic cell lines.

Images



Western blot analysis of LEF1 expression in Jurkat cell lysate.



Immunohistochemical analysis of paraffin-embedded human spleen, using LEF1 Antibody.