

Anti-M6PR IGF2R Rabbit Monoclonal Antibody

Catalog # ABO13381

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P11717
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-M6PR IGF2R Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	3482
Other Names	Cation-independent mannose-6-phosphate receptor, CI Man-6-P receptor, CI-MPR, M6PR, 300 kDa mannose 6-phosphate receptor, MPR 300, Insulin-like growth factor 2 receptor, Insulin-like growth factor II receptor, IGF-II receptor, M6P/IGF2 receptor, M6P/IGF2R, CD222, IGF2R, MPRI
Calculated MW	274375
Application Details	WB 1:5000-1:20000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:30 FC 1:100
Subcellular Localization	Lysosome membrane ; Single-pass type I membrane protein. Colocalized with DPP4 in internalized cytoplasmic vesicles adjacent to the cell surface.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: DIC-9
Immunogen	A synthesized peptide derived from human M6PR
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name IGF2R

Synonyms MPRI


Function Mediates the transport of phosphorylated lysosomal enzymes from the Golgi complex and the cell surface to lysosomes (PubMed:[18817523](#), PubMed:[2963003](#)). Lysosomal enzymes bearing phosphomannosyl residues bind specifically to mannose-6-phosphate receptors in the Golgi apparatus and the resulting receptor-ligand complex is transported to an acidic prelysosomal compartment where the low pH mediates the dissociation of the complex (PubMed:[18817523](#), PubMed:[2963003](#)). The receptor is then recycled back to the Golgi for another round of trafficking through its binding to the retromer (PubMed:[18817523](#)). This receptor also binds IGF2 (PubMed:[18046459](#)). Acts as a positive regulator of T-cell coactivation by binding DPP4 (PubMed:[10900005](#)).

Cellular Location Golgi apparatus membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Note=Mainly localized in the Golgi at steady state and not detectable in lysosome (PubMed:[18817523](#)) Colocalized with DPP4 in internalized cytoplasmic vesicles adjacent to the cell surface (PubMed:[10900005](#)).

Images



Western blot analysis of extracts of M6PR/IGF2R expression in Jurkat cell lysate.



Immunohistochemical analysis of paraffin-embedded human colon, using M6PR/IGF2R Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.