

Anti-GRK2 ADRBK1 Rabbit Monoclonal Antibody

Catalog # ABO13397

Product Information

Application	WB, IHC
Primary Accession	P25098
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-GRK2 ADRBK1 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	156
Other Names	Beta-adrenergic receptor kinase 1, Beta-ARK-1, 2.7.11.15, G-protein coupled receptor kinase 2 {ECO:0000312 HGNC:HGNC:289}, GRK2 (HGNC:289), ADRBK1, BARK, BARK1
Calculated MW	79574
Application Details	WB 1:500-1:2000 IHC 1:50-1:200
Subcellular Localization	Cytoplasm. Cell membrane.
Tissue Specificity	Expressed in peripheral blood leukocytes..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: EGE-1
Immunogen	A synthesized peptide derived from human GRK2
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	GRK2 {ECO:0000303 PubMed:40384633, ECO:0000312 HGNC:HGNC:289}
-------------	---

Function	Specifically phosphorylates the activated forms of G protein- coupled receptors (PubMed: 14976224 , PubMed: 16179383 , PubMed: 19306925 , PubMed: 19715378 , PubMed: 21799898 , PubMed: 23077507 , PubMed: 30284927 , PubMed: 39138140 , PubMed: 40384633). Such receptor phosphorylation initiates beta-arrestin-mediated receptor internalization, and signaling events leading to their desensitization (PubMed: 14976224 , PubMed: 16179383 , PubMed: 19306925 , PubMed: 19715378 , PubMed: 21799898 , PubMed: 23077507 , PubMed: 30284927 , PubMed: 39138140 , PubMed: 40384633). Phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors, probably inducing a desensitization of them (PubMed: 19715378). Phosphorylates catecholamine-activated ADRB2 to regulate physiological cardiomyocyte contraction rate responses (By similarity). Also phosphorylates ligand-bound C3a and C5a anaphylatoxin receptors (C3AR1 and C5AR1, respectively), leading to receptor desensitization (PubMed: 21799898 , PubMed: 23077507). Positively regulates ciliary smoothened (SMO) signaling pathway by mediating phosphorylation of SMO, facilitating SMO trafficking into the cilium and the stimulation of SMO activity (PubMed: 39138140). Also positively regulates SMO signaling by mediating phosphorylation of GPR161, a key inhibitor SMO signaling, promoting ciliary export of GPR161 (PubMed: 40384633). Key regulator of LPAR1 signaling (PubMed: 19306925). Competes with RALA for binding to LPAR1 thus affecting the signaling properties of the receptor (PubMed: 19306925). Desensitizes LPAR1 and LPAR2 in a phosphorylation-independent manner (PubMed: 19306925). Inhibits relaxation of airway smooth muscle in response to blue light (PubMed: 30284927).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:P26817}. Cell membrane {ECO:0000250 UniProtKB:P21146}. Cell projection, cilium membrane. Postsynapse {ECO:0000250 UniProtKB:P26817}. Presynapse {ECO:0000250 UniProtKB:P26817}
Tissue Location	Expressed in peripheral blood leukocytes.

Images



Western blot analysis of GRK2 expression in THP-1 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.