10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-SHP2 PTPN11 Rabbit Monoclonal Antibody

Catalog # ABO13432

Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	Q06124
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-SHP2 PTPN11 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human.


Additional Information

Gene ID	5781
Other Names	Tyrosine-protein phosphatase non-receptor type 11, 3.1.3.48, Protein-tyrosine phosphatase 1D, PTP-1D, Protein-tyrosine phosphatase 2C, PTP-2C, SH-PTP2, SHP-2, Shp2, SH-PTP3, PTPN11, PTP2C, SHPTP2
Calculated MW	68011
Application Details	WB 1:1000-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50
Subcellular Localization	Cytoplasm.
Tissue Specificity	Widely expressed, with highest levels in heart, brain, and skeletal muscle..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: GCA-16
Immunogen	A synthesized peptide derived from human SHP2
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Name	PTPN11
Synonyms	PTP2C, SHPTP2
Function	Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus (PubMed:10655584, PubMed:14739280, PubMed:18559669, PubMed:18829466, PubMed:26742426, PubMed:28074573, PubMed:32184441). Positively regulates MAPK signal transduction pathway (PubMed:28074573). Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed:28074573). Dephosphorylates ROCK2 at 'Tyr-722' resulting in stimulation of its RhoA binding activity (PubMed:18559669). Dephosphorylates CDC73 (PubMed:26742426). Dephosphorylates SOX9 on tyrosine residues, leading to inactivate SOX9 and promote ossification (By similarity). Dephosphorylates tyrosine-phosphorylated NEDD9/CAS-L (PubMed:19275884). Acts as an effector of PDCD1-mediated inhibition of T-cell response: recruited by phosphorylated PDCD1, mediating dephosphorylation of key T-cell receptor (TCR) proximal signaling molecules, leading to TCR signaling inhibition (PubMed:32184441).
Cellular Location	Cytoplasm. Nucleus
Tissue Location	Widely expressed, with highest levels in heart, brain, and skeletal muscle.

Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of SHP2 expression in Jurkat cell lysate.

Immunohistochemical analysis of paraffin-embedded human heart, using SHP2 Antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.