

Anti-IRF5 Rabbit Monoclonal Antibody

Catalog # ABO13446

Product Information

Application	WB, IHC, FC
Primary Accession	Q13568
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-IRF5 Rabbit Monoclonal Antibody . Tested in WB, IHC, Flow Cytometry applications. This antibody reacts with Human.

Additional Information

Gene ID	3663
Other Names	Interferon regulatory factor 5, IRF-5, IRF5 {ECO:0000303 PubMed:11303025, ECO:0000312 HGNC:HGNC:6120}
Calculated MW	56044
Application Details	WB 1:1000-1:2000 IHC 1:50-1:200 FC 1:100
Subcellular Localization	Cytoplasm. Nucleus. Shuttles between the nucleus and the cytoplasm.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: GGD-9
Immunogen	A synthesized peptide derived from human IRF5
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	IRF5 {ECO:0000303 PubMed:11303025, ECO:0000312 HGNC:HGNC:6120}
Function	Transcription factor that plays a critical role in innate immunity by activating

expression of type I interferon (IFN) IFNA and INFB and inflammatory cytokines downstream of endolysosomal toll-like receptors TLR7, TLR8 and TLR9 (PubMed:[11303025](#), PubMed:[15695821](#), PubMed:[22412986](#), PubMed:[25326418](#), PubMed:[32433612](#), PubMed:[40639948](#)). Regulates the transcription of type I IFN genes (IFN-alpha and IFN- beta) and IFN-stimulated genes (ISG) by binding to an interferon- stimulated response element (ISRE) in their promoters (By similarity). Can efficiently activate both the IFN-beta (IFNB) and the IFN-alpha (IFNA) genes and mediate their induction downstream of the TLR- activated, MyD88-dependent pathway (By similarity). Key transcription factor regulating the IFN response during SARS-CoV-2 infection (PubMed:[33440148](#)).


Cellular Location

Cytoplasm. Nucleus. Note=Shuttles between the nucleus and the cytoplasm: upon activation by the TLR adapter MYD88 and subsequent phosphorylation, translocates to the nucleus

Images



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of IRF5 expression in Ramos cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.