

Anti-PAX6 Rabbit Monoclonal Antibody

Catalog # ABO13470

Product Information

Application	WB, IHC, FC
Primary Accession	P26367
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-PAX6 Rabbit Monoclonal Antibody . Tested in WB, IHC, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	5080
Other Names	Paired box protein Pax-6, Aniridia type II protein, Oculorhombin, PAX6, AN2
Calculated MW	46683
Application Details	WB 1:1000-1:2000 IHC 1:50-1:200 FC 1:500
Subcellular Localization	Nucleus.
Tissue Specificity	Fetal eye, brain, spinal cord and olfactory epithelium. Isoform 5a is less abundant than the PAX6 shorter form.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: HEB-16
Immunogen	A synthesized peptide derived from human PAX6
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	PAX6
-------------	------

Synonyms	AN2
Function	Transcription factor with important functions in the development of the eye, nose, central nervous system and pancreas. Required for the differentiation of pancreatic islet alpha cells (By similarity). Competes with PAX4 in binding to a common element in the glucagon, insulin and somatostatin promoters. Regulates specification of the ventral neuron subtypes by establishing the correct progenitor domains (By similarity). Acts as a transcriptional repressor of NFATC1- mediated gene expression (By similarity).
Cellular Location	Nucleus {ECO:0000250 UniProtKB:P63015}. [Isoform 5a]: Nucleus {ECO:0000250 UniProtKB:P63016}
Tissue Location	[Isoform 1]: Expressed in lymphoblasts.


Images



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.

Western blot analysis of PAX6 expression in HeLa cell lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.