

Anti-CDC5 CDC5L Rabbit Monoclonal Antibody

Catalog # ABO13480

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	Q99459
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CDC5 CDC5L Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Gene ID	988
Other Names	Cell division cycle 5-like protein, Cdc5-like protein, Pombe cdc5-related protein, CDC5L, KIAA0432, PCDC5RP
Calculated MW	92251
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Subcellular Localization	Nucleus. Nucleus speckle. Cytoplasm. May shuttle between cytoplasm and nucleus..
Tissue Specificity	Ubiquitously expressed in both fetal and adult tissues..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: IOA-3
Immunogen	A synthesized peptide derived from human CDC5
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CDC5L
Synonyms	KIAA0432, PCDC5RP
Function	DNA-binding protein involved in cell cycle control. May act as a transcription activator. Plays a role in pre-mRNA splicing as core component of precatalytic, catalytic and postcatalytic spliceosomal complexes (PubMed: 11991638 , PubMed: 20176811 , PubMed: 28076346 , PubMed: 28502770 , PubMed: 29301961 , PubMed: 29360106 , PubMed: 29361316 , PubMed: 30705154 , PubMed: 30728453). Component of the PRP19-CDC5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing. The PRP19-CDC5L complex may also play a role in the response to DNA damage (DDR) (PubMed: 20176811). As a component of the minor spliceosome, involved in the splicing of U12- type introns in pre-mRNAs (Probable).
Cellular Location	Nucleus. Nucleus speckle. Cytoplasm Note=May shuttle between cytoplasm and nucleus
Tissue Location	Ubiquitously expressed in both fetal and adult tissues.

Images



Western blot analysis of CDC5 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.