

Anti-E2F1 Rabbit Monoclonal Antibody

Catalog # ABO13491

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	Q01094
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-E2F1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Gene ID	1869
Other Names	Transcription factor E2F1, E2F-1, PBR3, Retinoblastoma-associated protein 1, RBAP-1, Retinoblastoma-binding protein 3, RBBP-3, pRB-binding protein E2F-1, E2F1 {ECO:0000303 PubMed:8964493, ECO:0000312 HGNC:HGNC:3113}
Calculated MW	46920
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Subcellular Localization	Nucleus.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: IGD-5
Immunogen	A synthesized peptide derived from human E2F1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Name	E2F1 {ECO:0000303 PubMed:8964493, ECO:0000312 HGNC:HGNC:3113}
Function	Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication (PubMed: 10675335 , PubMed: 12717439 , PubMed: 17050006 , PubMed: 17704056 , PubMed: 18625225 , PubMed: 28992046). The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase (PubMed: 10675335 , PubMed: 12717439 , PubMed: 17704056). E2F1 binds preferentially RB1 in a cell-cycle dependent manner (PubMed: 10675335 , PubMed: 12717439 , PubMed: 17704056). It can mediate both cell proliferation and TP53/p53-dependent apoptosis (PubMed: 8170954). Blocks adipocyte differentiation by binding to specific promoters repressing CEBPA binding to its target gene promoters (PubMed: 20176812). Directly activates transcription of PEG10 (PubMed: 17050006 , PubMed: 18625225 , PubMed: 28992046). Positively regulates transcription of RRP1B (PubMed: 20040599).
Cellular Location	Nucleus


Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




Immunofluorescent analysis using the Antibody at 1:50 dilution.




Immunofluorescent analysis using the Antibody at 1:50 dilution.



Western blot analysis of E2F1 expression in (1) HeLa cell lysate; (2) NIH/3T3 cell lysate.



Immunohistochemical analysis of paraffin-embedded human ovary carcinoma, using E2F1 Antibody.



Immunofluorescent analysis of HeLa cells, using E2F1 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.