

Anti-CD19 Rabbit Monoclonal Antibody

Catalog # ABO13504

Product Information

Application	WB, FC
Primary Accession	P15391
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CD19 Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	930
Other Names	B-lymphocyte antigen CD19, B-lymphocyte surface antigen B4, Differentiation antigen CD19, T-cell surface antigen Leu-12, CD19, CD19
Calculated MW	61128
Application Details	WB 1:500-1:1000 FC 1:50
Subcellular Localization	Membrane; Single-pass type I membrane protein.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AOEB-3
Immunogen	A synthesized peptide derived from human CD19
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CD19
Function	Functions as a coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes (PubMed: 29523808). Decreases the threshold for activation of

downstream signaling pathways and for triggering B-cell responses to antigens (PubMed:[1373518](#), PubMed:[16672701](#), PubMed:[2463100](#)). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed:[12387743](#), PubMed:[16672701](#), PubMed:[9317126](#), PubMed:[9382888](#)). Is not required for early steps during B cell differentiation in the blood marrow (PubMed:[9317126](#)). Required for normal differentiation of B-1 cells (By similarity). Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed:[1373518](#), PubMed:[2463100](#)). Required for normal levels of serum immunoglobulins, and for production of high-affinity antibodies in response to antigen challenge (PubMed:[12387743](#), PubMed:[16672701](#), PubMed:[9317126](#)).


Cellular Location

Cell membrane; Single-pass type I membrane protein. Membrane raft {ECO:0000250|UniProtKB:P25918}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P25918}


Tissue Location

Detected on marginal zone and germinal center B cells in lymph nodes (PubMed:[2463100](#)). Detected on blood B cells (at protein level) (PubMed:[16672701](#), PubMed:[2463100](#))

Images




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.

Western blot analysis of CD19 expression in (1) Jurkat cell lysate; (2) Rat spleen lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.