

Anti-IP10 CXCL10 Rabbit Monoclonal Antibody

Catalog # ABO13530

Product Information

Application	WB
Primary Accession	P02778
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-IP10 CXCL10 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.

Additional Information


Gene ID	3627
Other Names	C-X-C motif chemokine 10, 10 kDa interferon gamma-induced protein, Gamma-IP10, IP-10, Small-inducible cytokine B10, CXCL10(1-73), CXCL10, INP10, SCYB10
Calculated MW	10881
Application Details	WB 1:500-1:2000
Subcellular Localization	Secreted.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AACI-3
Immunogen	A synthesized peptide derived from human IP10
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CXCL10
Synonyms	INP10, SCYB10

Function	Pro-inflammatory cytokine that is involved in a wide variety of processes such as chemotaxis, differentiation, and activation of peripheral immune cells, regulation of cell growth, apoptosis and modulation of angiostatic effects (PubMed: 11157474 , PubMed: 22652417 , PubMed: 7540647). Plays thereby an important role during viral infections by stimulating the activation and migration of immune cells to the infected sites (By similarity). Mechanistically, binding of CXCL10 to the CXCR3 receptor activates G protein-mediated signaling and results in downstream activation of phospholipase C-dependent pathway, an increase in intracellular calcium production and actin reorganization (PubMed: 12750173 , PubMed: 19151743). In turn, recruitment of activated Th1 lymphocytes occurs at sites of inflammation (PubMed: 12663757 , PubMed: 12750173). Activation of the CXCL10/CXCR3 axis also plays an important role in neurons in response to brain injury for activating microglia, the resident macrophage population of the central nervous system, and directing them to the lesion site. This recruitment is an essential element for neuronal reorganization (By similarity).
Cellular Location	Secreted.
Tissue Location	Mainly secreted by monocytes, endothelial cells as well as fibroblasts. Expressed by epithelial cells in thymus (PubMed: 11157474). Microglial cells produce CXCL10 in response to viral stimulation (PubMed: 12663757).

Images



Western blot analysis of IP10 expression in IP10 recombinant protein lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.