

Anti-AQP1 Rabbit Monoclonal Antibody

Catalog # ABO13536

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P29972
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-AQP1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	358
Other Names	Aquaporin-1, AQP-1, Aquaporin-CHIP, Urine water channel, Water channel protein for red blood cells and kidney proximal tubule, AQP1 (HGNC:633), CHIP28
Calculated MW	28526
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Subcellular Localization	Cell membrane ; Multi-pass membrane protein.
Tissue Specificity	Detected in erythrocytes (at protein level). Expressed in a number of tissues including erythrocytes, renal tubules, retinal pigment epithelium, heart, lung, skeletal muscle, kidney and pancreas. Weakly expressed in brain, placenta and liver..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AAEE-1
Immunogen	A synthesized peptide derived from human AQP1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name AQP1 ([HGNC:633](#))

Function Forms a water channel that facilitates the transport of water across cell membranes, playing a crucial role in water homeostasis in various tissues (PubMed:[1373524](#), PubMed:[23219802](#)). Could also be permeable to small solutes including hydrogen peroxide, glycerol and gases such as ammonia (NH₃), nitric oxide (NO) and carbon dioxide (CO₂) (PubMed:[16682607](#), PubMed:[17012249](#), PubMed:[19273840](#), PubMed:[33028705](#), PubMed:[8584435](#)). Recruited to the ankyrin-1 complex, a multiprotein complex of the erythrocyte membrane, it could be part of a CO₂ metabolon, linking facilitated diffusion of CO₂ across the membrane, anion exchange of Cl⁻/HCO₃⁻ and interconversion of dissolved CO₂ and carbonic acid in the cytosol (PubMed:[17012249](#), PubMed:[35835865](#)). In vitro, it shows non-selective gated cation channel activity and may be permeable to cations like K⁺ and Na⁺ in vivo (PubMed:[36949749](#), PubMed:[8703053](#)).


Cellular Location Cell membrane; Multi-pass membrane protein

Tissue Location Detected in erythrocytes (at protein level). Expressed in a number of tissues including erythrocytes, renal tubules, retinal pigment epithelium, heart, lung, skeletal muscle, kidney and pancreas. Weakly expressed in brain, placenta and liver

Images



Western blot analysis of AQP1 expression in (1) Human fetal kidney lysate; (2) Human fetal lung lysate.



Immunohistochemical analysis of paraffin-embedded human breast, using AQP1 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.