10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD18 ITGB2 Rabbit Monoclonal Antibody

Catalog # ABO13543

Product Information

Application	WB, IP
Primary Accession	P05107
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-CD18 ITGB2 Rabbit Monoclonal Antibody . Tested in WB, IP applications. This antibody reacts with Human.

Additional Information

Gene ID	3689
Other Names	Integrin beta-2, Cell surface adhesion glycoproteins LFA-1/CR3/p150, 95 subunit beta, Complement receptor C3 subunit beta, CD18, ITGB2, CD18, MFI7
Calculated MW	84791
Application Details	WB 1:500-1:2000 IP 1:50
Subcellular Localization	Membrane; Single-pass type I membrane protein.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AAIH-9
Immunogen	A synthesized peptide derived from human CD18
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	ITGB2 (HGNC:6155)
Synonyms	CD18, MFI7

Function

Integrin ITGAL:ITGB2 is a receptor for ICAM1, ICAM2 and ICAM3 (PubMed:1676048, PubMed:23775590, PubMed:38195629). Integrin ITGAL:ITGB2 is also a receptor for the secreted form of ubiquitin-like protein ISG15; the interaction is mediated by ITGAL (PubMed:29100055). Integrins ITGAM:ITGB2 and ITGAX:ITGB2 are receptors for the iC3b fragment of the third complement component and for fibrinogen. Integrin ITGAX:ITGB2 recognizes the sequence G-P-R in fibrinogen alpha-chain. Integrin ITGAM:ITGB2 recognizes P1 and P2 peptides of fibrinogen gamma chain. Integrin ITGAM:ITGB2 is also a receptor for factor X. Integrin ITGAD:ITGB2 is a receptor for ICAM3 and VCAM1 (PubMed:10438935, PubMed:8777714, PubMed:9841932). Contributes to natural killer cell cytotoxicity (PubMed:15356110). Involved in leukocyte adhesion and transmigration of leukocytes including T-cells and neutrophils (PubMed:11812992, PubMed:28807980). Triggers neutrophil transmigration during lung injury through PTK2B/PYK2-mediated activation (PubMed:18587400). Integrin ITGAL:ITGB2 in association with ICAM3, contributes to apoptotic neutrophil phagocytosis by macrophages (PubMed:23775590). In association with alpha subunit ITGAM/CD11b, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed:21193407). Integrins ITGAX:ITGB2 functions as a receptor of the erythrocyte-specific adhesion molecule ICAM4 and mediates erythrophagocytosis (PubMed:16985175). Integrins ITGAX:ITGB2 functions as a receptor of the neuron-specific adhesion molecule ICAM5 ensuring neuron cell-leukocyte adhesion (PubMed:10741396). Integrin ITGAL:ITGB2 functions as a receptor of ICAM1 by acting as a platform at the immunological synapse to translate TCR engagement and density of the ITGAL ligand ICAM1 into graded adhesion (PubMed:38195629). Integrin ITGAM:ITGB2/MAC-1 complex functions as a signaling receptor for the ligand receptor ICAM1, ensuring adhesion between stimulated neutrophils and stimulated endothelial cells (PubMed:1980124). Integrin ITGAL/ITGB2 that functions as a signaling receptor of ICAM2, ensuring leukocyte cell-cell adhesion on resting cells (PubMed:1676048).


Cellular Location

Cell membrane; Single-pass type I membrane protein. Membrane raft; Single-pass type I membrane protein

Tissue Location

Leukocytes (PubMed:23775590). Expressed in neutrophils (at protein level) (PubMed:21193407, PubMed:28807980)

Images



Western blot analysis of CD18 expression in K562 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.