

Anti-PAI1 Rabbit Monoclonal Antibody

Catalog # ABO13564

Product Information

Application	WB, IF, ICC, IP, FC
Primary Accession	P05121
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-PAI1 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.


Additional Information

Gene ID	5054
Other Names	Plasminogen activator inhibitor 1, PAI, PAI-1, Endothelial plasminogen activator inhibitor, Serpin E1, SERPINE1, PAI1, PLANH1
Calculated MW	45060
Application Details	WB 1:500-1:1000 ICC/IF 1:50-1:200 IP 1:30 FC 1:50
Subcellular Localization	Secreted.
Tissue Specificity	Found in plasma and platelets and in endothelial, hepatoma and fibrosarcoma cells.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ABHG-19
Immunogen	A synthesized peptide derived from human PAI1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	SERPINE1
Synonyms	PAI1, PLANH1
Function	Serine protease inhibitor. Inhibits TMPRSS7 (PubMed: 15853774). Is a primary inhibitor of tissue-type plasminogen activator (PLAT) and urokinase-type plasminogen activator (PLAU). As PLAT inhibitor, it is required for fibrinolysis down-regulation and is responsible for the controlled degradation of blood clots (PubMed: 17912461 , PubMed: 8481516 , PubMed: 9207454 , PubMed: 21925150). As PLAU inhibitor, it is involved in the regulation of cell adhesion and spreading (PubMed: 9175705). Acts as a regulator of cell migration, independently of its role as protease inhibitor (PubMed: 15001579 , PubMed: 9168821). It is required for stimulation of keratinocyte migration during cutaneous injury repair (PubMed: 18386027). It is involved in cellular and replicative senescence (PubMed: 16862142). Plays a role in alveolar type 2 cells senescence in the lung (By similarity). Is involved in the regulation of cementogenic differentiation of periodontal ligament stem cells, and regulates odontoblast differentiation and dentin formation during odontogenesis (PubMed: 25808697 , PubMed: 27046084).
Cellular Location	Secreted.
Tissue Location	Expressed in endothelial cells (PubMed:2430793, PubMed:3097076). Found in plasma, platelets, and hepatoma and fibrosarcoma cells.

Images



Western blot analysis of PAI1 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.