

Anti-MMP14/Mt1 Mmp Rabbit Monoclonal Antibody

Catalog # ABO13608

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P50281
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-MMP14/Mt1 Mmp Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	4323
Other Names	Matrix metalloproteinase-14, MMP-14, 3.4.24.80, MMP-X1, Membrane-type matrix metalloproteinase 1, MT-MMP 1, MTMMP1, Membrane-type-1 matrix metalloproteinase, MT1-MMP, MT1MMP, MMP14
Calculated MW	65894
Application Details	WB 1:1000-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Subcellular Localization	Membrane ; Single-pass type I membrane protein. Melanosome. Cytoplasm. Identified by mass spectrometry in melanosome fractions from stage I to stage IV. Forms a complex with BST2 and localizes to the cytoplasm.
Tissue Specificity	Expressed in stromal cells of colon, breast, and head and neck. Expressed in lung tumors..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: BFD-13
Immunogen	A synthesized peptide derived from human MMP14
Purification	Affinity-chromatography

Storage Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name MMP14

Function Endopeptidase that degrades various components of the extracellular matrix such as collagen (PubMed:[8015608](#)). Essential for pericellular collagenolysis and modeling of skeletal and extraskelatal connective tissues during development (By similarity). Activates progelatinase A/MMP2, thereby acting as a positive regulator of cell growth and migration (PubMed:[22065321](#), PubMed:[8015608](#)). Involved in the formation of the fibrovascular tissues in association with pro-MMP2 (PubMed:[12714657](#), PubMed:[22065321](#)). May be involved in actin cytoskeleton reorganization by cleaving PTK7 (PubMed:[20837484](#)). Acts as a regulator of Notch signaling by mediating cleavage and inhibition of DLL1 (PubMed:[21572390](#)). Cleaves ADGRB1 to release vasculostatin-40 which inhibits angiogenesis (PubMed:[22330140](#)). Acts as a negative regulator of the GDF15-GFRAL aversive response by mediating cleavage and inactivation of GFRAL (PubMed:[35177851](#)).


Cellular Location Cell membrane; Single-pass type I membrane protein. Melanosome. Cytoplasm Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:[17081065](#)). Forms a complex with BST2 and localizes to the cytoplasm (PubMed:[17081065](#))

Tissue Location Expressed in stromal cells of colon, breast, and head and neck. Expressed in lung tumors.

Images



Western blot analysis of MMP14 expression in human spleen lysate.



Immunohistochemical analysis of paraffin-embedded human kidney, using MMP14 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.