

Anti-PSD95 DLG4 Rabbit Monoclonal Antibody

Catalog # ABO13633

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P78352
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-PSD95 DLG4 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	1742
Other Names	Disks large homolog 4, Postsynaptic density protein 95, PSD-95, Synapse-associated protein 90, SAP-90, SAP90, DLG4 (HGNC:2903), PSD95
Calculated MW	80495
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:20 FC 1:20
Subcellular Localization	Cell membrane ; Peripheral membrane protein. Cell junction, synapse, postsynaptic cell membrane, postsynaptic density. Cell projection, axon. Cell junction, synapse. High levels in postsynaptic density of neurons in the forebrain. Also in presynaptic region of inhibitory synapses formed by cerebellar basket cells on axon hillocks of Purkinje cells.
Tissue Specificity	Brain.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: DAF-4
Immunogen	A synthesized peptide derived from human PSD95
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at

4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	DLG4 (HGNC:2903)
Synonyms	PSD95
Function	Postsynaptic scaffolding protein that plays a critical role in synaptogenesis and synaptic plasticity by providing a platform for the postsynaptic clustering of crucial synaptic proteins. Interacts with the cytoplasmic tail of NMDA receptor subunits and shaker-type potassium channels. Required for synaptic plasticity associated with NMDA receptor signaling. Overexpression or depletion of DLG4 changes the ratio of excitatory to inhibitory synapses in hippocampal neurons. May reduce the amplitude of ASIC3 acid-evoked currents by retaining the channel intracellularly. May regulate the intracellular trafficking of ADR1B. Also regulates AMPA-type glutamate receptor (AMPA) immobilization at postsynaptic density keeping the channels in an activated state in the presence of glutamate and preventing synaptic depression (By similarity). Under basal conditions, cooperates with FYN to stabilize palmitoyltransferase ZDHHC5 at the synaptic membrane through FYN-mediated phosphorylation of ZDHHC5 and its subsequent inhibition of association with endocytic proteins (PubMed: 26334723).
Cellular Location	Cell membrane; Lipid-anchor; Cytoplasmic side. Postsynaptic density {ECO:0000250 UniProtKB:P31016}. Synapse Cytoplasm {ECO:0000250 UniProtKB:P31016}. Cell projection, axon {ECO:0000250 UniProtKB:P31016}. Cell projection, dendritic spine {ECO:0000250 UniProtKB:P31016}. Cell projection, dendrite {ECO:0000250 UniProtKB:P31016}. Presynapse {ECO:0000250 UniProtKB:P31016}. Note=High levels in postsynaptic density of neurons in the forebrain. Also in presynaptic region of inhibitory synapses formed by cerebellar basket cells on axon hillocks of Purkinje cells. Suppression of neuronal activity induces synaptic accumulation and clustering of DLG4. {ECO:0000250 UniProtKB:P31016}
Tissue Location	Brain.

Images



Immunofluorescent analysis of U87-MG cells, using PSD95 Antibody .

Figure 1. Western blot analysis of DLG4 using anti-DLG4 antibody (M02128). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours.



The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat brain tissue lysates,

Lane 2: rat C6 whole cell lysates,

Lane 3: mouse brain tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-DLG4 antigen affinity purified monoclonal antibody (Catalog # M02128) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for DLG4 at approximately 95 kDa. The expected band size for DLG4 is at 80 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.