

# Anti-SOX18 Rabbit Monoclonal Antibody

Catalog # ABO13638

## Product Information

---

|                          |                                                                                                          |
|--------------------------|----------------------------------------------------------------------------------------------------------|
| <b>Application</b>       | WB, IHC                                                                                                  |
| <b>Primary Accession</b> | <a href="#">P35713</a>                                                                                   |
| <b>Host</b>              | Rabbit                                                                                                   |
| <b>Isotype</b>           | Rabbit IgG                                                                                               |
| <b>Reactivity</b>        | Human                                                                                                    |
| <b>Clonality</b>         | Monoclonal                                                                                               |
| <b>Format</b>            | Liquid                                                                                                   |
| <b>Description</b>       | Anti-SOX18 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human. |

## Additional Information

---

|                                 |                                                                                                                                            |
|---------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Gene ID</b>                  | 54345                                                                                                                                      |
| <b>Other Names</b>              | Transcription factor SOX-18, SOX18                                                                                                         |
| <b>Calculated MW</b>            | 40891                                                                                                                                      |
| <b>Application Details</b>      | WB 1:500-1:2000<br>IHC 1:50-1:200                                                                                                          |
| <b>Subcellular Localization</b> | Nucleus.                                                                                                                                   |
| <b>Source</b>                   | Eukaryota                                                                                                                                  |
| <b>Contents</b>                 | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.                        |
| <b>Clone Names</b>              | Clone: DCD-19                                                                                                                              |
| <b>Immunogen</b>                | A synthesized peptide derived from human SOX18                                                                                             |
| <b>Purification</b>             | Affinity-chromatography                                                                                                                    |
| <b>Storage</b>                  | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles. |

## Protein Information

---

|                 |                                                                                                                                                                                                                   |
|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Name</b>     | SOX18                                                                                                                                                                                                             |
| <b>Function</b> | Transcriptional activator that binds to the consensus sequence 5'-AACAAAG-3' in the promoter of target genes and plays an essential role in embryonic cardiovascular development and lymphangiogenesis. Activates |

transcription of PROX1 and other genes coding for lymphatic endothelial markers. Plays an essential role in triggering the differentiation of lymph vessels, but is not required for the maintenance of differentiated lymphatic endothelial cells. Plays an important role in postnatal angiogenesis, where it is functionally redundant with SOX17. Interaction with MEF2C enhances transcriptional activation. Besides, required for normal hair development.

#### Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00267}.

#### Tissue Location

Detected in heart, lung, placenta, skeletal muscle, liver, kidney, spleen, prostate, ovary, msosmall intestine and colon

## Images




Figure 1. Western blot analysis of SOX18 using anti-SOX18 antibody (M04004).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human K562 whole cell lysates,

Lane 2: human COLO 320 whole cell lysates,

Lane 3: human HepG2 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SOX18 antigen affinity purified monoclonal antibody (Catalog # M04004) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SOX18 at approximately 41 kDa. The expected band size for SOX18 is at 41 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.