

Anti-NEDD8 Rabbit Monoclonal Antibody

Catalog # ABO13732

Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	Q15843
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-NEDD8 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Gene ID	4738
Other Names	NEDD8, Neddylin, Neural precursor cell expressed developmentally down-regulated protein 8, NEDD-8, Ubiquitin-like protein Nedd8, NEDD8 {ECO:0000303 PubMed:9694792, ECO:0000312 HGNC:HGNC:7732}
Calculated MW	9072
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:20
Subcellular Localization	Nucleus. Mainly nuclear.
Tissue Specificity	Highly expressed in heart, skeletal muscle, spleen, thymus, prostate, testis, ovary, colon and leukocytes..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AOA-E-14
Immunogen	A synthesized peptide derived from human NEDD8
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	NEDD8 {ECO:0000303 PubMed:9694792, ECO:0000312 HGNC:HGNC:7732}
Function	Ubiquitin-like protein which plays an important role in cell cycle control and embryogenesis via its conjugation to a limited number of cellular proteins, such as cullins or p53/TP53 (PubMed: 10318914 , PubMed: 10597293 , PubMed: 11953428 , PubMed: 14690597 , PubMed: 15242646 , PubMed: 9694792 , PubMed: 38605244 , PubMed: 38316879). Attachment of NEDD8 to cullins is critical for the recruitment of E2 to the cullin-RING- based E3 ubiquitin-protein ligase complex, thus facilitating polyubiquitination and proteasomal degradation of cyclins and other regulatory proteins (PubMed: 10318914 , PubMed: 10597293 , PubMed: 11953428 , PubMed: 20688984 , PubMed: 9694792 , PubMed: 38605244 , PubMed: 38316879). Attachment of NEDD8 to p53/TP53 inhibits p53/TP53 transcriptional activity (PubMed: 15242646). Covalent attachment to its substrates requires prior activation by the E1 complex UBE1C-APPBP1 and linkage to the E2 enzyme UBE2M (PubMed: 14690597). Neddylates EGFR resulting in protein stabilization and translocation to the cell membrane (PubMed: 28891468).
Cellular Location	Nucleus. Note=Mainly nuclear.
Tissue Location	Highly expressed in heart, skeletal muscle, spleen, thymus, prostate, testis, ovary, colon and leukocytes

Images



Western blot analysis of NEDD8 expression in K562 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.