

Anti-PDIA6/Erp5 Rabbit Monoclonal Antibody

Catalog # ABO13735

Product Information

Application	WB, IF, ICC, FC
Primary Accession	Q15084
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-PDIA6/Erp5 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Gene ID	10130
Other Names	Protein disulfide-isomerase A6, 5.3.4.1, Endoplasmic reticulum protein 5, ER protein 5, ERp5, Protein disulfide isomerase P5, Thioredoxin domain-containing protein 7, PDIA6, ERP5, P5, TXNDC7
Calculated MW	48121
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 FC 1:50
Subcellular Localization	Endoplasmic reticulum lumen. Cell membrane. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545)..
Tissue Specificity	Expressed in platelets (at protein level)..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AOB6-16
Immunogen	A synthesized peptide derived from human PDIA6
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	PDIA6
Synonyms	ERP5, P5, TXNDC7
Function	May function as a chaperone that inhibits aggregation of misfolded proteins (PubMed: 12204115). Negatively regulates the unfolded protein response (UPR) through binding to UPR sensors such as ERN1, which in turn inactivates ERN1 signaling (PubMed: 24508390). May also regulate the UPR via the EIF2AK3 UPR sensor (PubMed: 24508390). Plays a role in platelet aggregation and activation by agonists such as convulxin, collagen and thrombin (PubMed: 15466936).
Cellular Location	Endoplasmic reticulum lumen. Cell membrane. Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545)
Tissue Location	Expressed in platelets (at protein level).

Images



Western blot analysis of PDIA6 expression in K562 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.