

Anti-CD147 BSG Rabbit Monoclonal Antibody

Catalog # ABO13753

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P35613
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CD147 BSG Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	682
Other Names	Basigin, 5F7, Collagenase stimulatory factor, Extracellular matrix metalloproteinase inducer, EMMPRIN, Hepatoma-associated antigen, HAB18G, Leukocyte activation antigen M6, OK blood group antigen, Tumor cell-derived collagenase stimulatory factor, TCSF, CD147, BSG (HGNC:1116)
Calculated MW	42200
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Subcellular Localization	Cell membrane ; Single-pass type I membrane protein. Melanosome. Colocalizes with SLC16A1 and SLC16A8. Identified by mass spectrometry in melanosome fractions from stage I to stage IV..
Tissue Specificity	Present only in vascular endothelium in non- neoplastic regions of the brain, whereas it is present in tumor cells but not in proliferating blood vessels in malignant gliomas.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AAOC-2
Immunogen	A synthesized peptide derived from human CD147
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at

4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Name	BSG (HGNC:1116)
Function	[Isoform 1]: Essential for normal retinal maturation and development (By similarity). Acts as a retinal cell surface receptor for NXNL1 and plays an important role in NXNL1-mediated survival of retinal cone photoreceptors (PubMed: 25957687). In association with glucose transporter SLC16A1/GLUT1 and NXNL1, promotes retinal cone survival by enhancing aerobic glycolysis and accelerating the entry of glucose into photoreceptors (PubMed: 25957687). May act as a potent stimulator of IL6 secretion in multiple cell lines that include monocytes (PubMed: 21620857).
Cellular Location	Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. [Isoform 2]: Cell membrane; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P26453}. Endosome Endoplasmic reticulum membrane; Single- pass type I membrane protein {ECO:0000250 UniProtKB:P26453} Basolateral cell membrane; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P26453} [Isoform 4]: Cell membrane; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P26453}
Tissue Location	[Isoform 1]: Retina-specific (PubMed:25957687). Expressed in retinal cone photoreceptors (at protein level) (PubMed:25957687). [Isoform 3]: Highly expressed in the bone marrow, fetal liver, lung, testis and thymus.

Images




Western blot analysis of CD147 expression in Mouse kidney lysate.



Immunohistochemical analysis of paraffin-embedded human stomach cancer, using CD147 Antibody.

Immunofluorescent analysis of Jurkat cells, using using CD147 Antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.