

Anti-ERp57 PDIA3 Rabbit Monoclonal Antibody

Catalog # ABO13782

Product Information

Application	WB, IHC, IP
Primary Accession	P30101
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-ERp57 PDIA3 Rabbit Monoclonal Antibody . Tested in WB, IHC, IP applications. This antibody reacts with Human.


Additional Information

Gene ID	2923
Other Names	Protein disulfide-isomerase A3, 5.3.4.1, 58 kDa glucose-regulated protein, 58 kDa microsomal protein, p58, Disulfide isomerase ER-60, Endoplasmic reticulum resident protein 57, ER protein 57, ERp57, Endoplasmic reticulum resident protein 60, ER protein 60, ERp60, PDIA3 (HGNC:4606), ERP57, ERP60, GRP58
Calculated MW	56782
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 IP 1:50
Subcellular Localization	Endoplasmic reticulum. Endoplasmic reticulum lumen. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.
Tissue Specificity	Detected in the flagellum and head region of spermatozoa (at protein level)..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ABOE-16
Immunogen	A synthesized peptide derived from human ERp57
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Name	PDIA3 (HGNC:4606)
Synonyms	ERP57, ERP60, GRP58
Function	Protein disulfide isomerase that catalyzes the formation, isomerization, and reduction or oxidation of disulfide bonds in client proteins and functions as a protein folding chaperone (PubMed: 11825568 , PubMed: 16193070 , PubMed: 27897272 , PubMed: 36104323 , PubMed: 7487104). Core component of the major histocompatibility complex class I (MHC I) peptide loading complex where it functions as an essential folding chaperone for TAPBP. Through TAPBP, assists the dynamic assembly of the MHC I complex with high affinity antigens in the endoplasmic reticulum. Therefore, plays a crucial role in the presentation of antigens to cytotoxic T cells in adaptive immunity (PubMed: 35948544 , PubMed: 36104323).
Cellular Location	Endoplasmic reticulum. Endoplasmic reticulum lumen {ECO:0000250 UniProtKB:P11598}. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545).
Tissue Location	Detected in the flagellum and head region of spermatozoa (at protein level) (PubMed:20400973). Expressed in liver, stomach and colon (at protein level). Expressed in gastric parietal cells and chief cells (at protein level) (PubMed:24188822)


Images



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



Western blot analysis of ERp57 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.