

Anti-BNIP1 Rabbit Monoclonal Antibody

Catalog # ABO13809

Product Information

Application	WB, IHC
Primary Accession	Q12981
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-BNIP1 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Gene ID	662
Other Names	Vesicle transport protein SEC20, BCL2/adenovirus E1B 19 kDa protein-interacting protein 1, Transformation-related gene 8 protein, TRG-8, BNIP1, NIP1, SEC20L
Calculated MW	26132
Application Details	WB 1:500-1:2000 IHC 1:50-1:200
Subcellular Localization	Mitochondrion. Endoplasmic reticulum membrane; Single-pass type IV membrane protein.
Tissue Specificity	Isoform 1 is highly expressed in heart, brain, liver skeletal muscle and pancreas. Isoform 3 is moderately expressed in placenta, lung and kidney. Isoform 4 is highly expressed in testis and small intestine..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ABHI-2
Immunogen	A synthesized peptide derived from human BNIP1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	BNIP1
Synonyms	NIP1, SEC20L
Function	As part of a SNARE complex may be involved in endoplasmic reticulum membranes fusion and be required for the maintenance of endoplasmic reticulum organization (PubMed: 15272311). Also plays a role in apoptosis (PubMed: 15272311 , PubMed: 23896122 , PubMed: 7954800). It is for instance required for endoplasmic reticulum stress-induced apoptosis (PubMed: 23896122). As a substrate of RNF185 interacting with SQSTM1, might also be involved in mitochondrial autophagy (Probable).
Cellular Location	Endoplasmic reticulum membrane; Single-pass type IV membrane protein. Mitochondrion membrane; Single-pass type IV membrane protein. Note=Localization to the mitochondrion is regulated by RNF186.
Tissue Location	Isoform 1 is highly expressed in heart, brain, liver skeletal muscle and pancreas. Isoform 3 is moderately expressed in placenta, lung and kidney. Isoform 4 is highly expressed in testis and small intestine.

Images



Western blot analysis of BNIP1 expression in (1) Jurkat cell lysate; (2) C6 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.