

Anti-Moesin Rabbit Monoclonal Antibody

Catalog # ABO13841

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P26038
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Moesin Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	4478
Other Names	Moesin, Membrane-organizing extension spike protein, MSN (HGNC:7373)
Calculated MW	67820
Application Details	WB 1:1000-1:2000 IHC 1:50-1:100 ICC/IF 1:50-1:100 IP 1:20 FC 1:20
Subcellular Localization	Cell membrane ; Peripheral membrane protein ; Cytoplasmic side. Cytoplasm, cytoskeleton. Apical cell membrane ; Peripheral membrane protein ; Cytoplasmic side. Cell projection, microvillus membrane ; Peripheral membrane protein ; Cytoplasmic side. Phosphorylated form is enriched in microvilli-like structures at apical membrane (By similarity). Increased cell membrane localization of both phosphorylated and non-phosphorylated forms seen after thrombin treatment..
Tissue Specificity	In all tissues and cultured cells studied.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: FIC-13
Immunogen	A synthesized peptide derived from human Moesin
Purification	Affinity-chromatography

Storage Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name MSN ([HGNC:7373](#))

Function Ezrin-radixin-moesin (ERM) family protein that connects the actin cytoskeleton to the plasma membrane and thereby regulates the structure and function of specific domains of the cell cortex. Tethers actin filaments by oscillating between a resting and an activated state providing transient interactions between moesin and the actin cytoskeleton (PubMed:[10212266](#)). Once phosphorylated on its C-terminal threonine, moesin is activated leading to interaction with F-actin and cytoskeletal rearrangement (PubMed:[10212266](#)). These rearrangements regulate many cellular processes, including cell shape determination, membrane transport, and signal transduction (PubMed:[12387735](#), PubMed:[15039356](#)). The role of moesin is particularly important in immunity acting on both T and B-cells homeostasis and self-tolerance, regulating lymphocyte egress from lymphoid organs (PubMed:[9298994](#), PubMed:[9616160](#)). Modulates phagolysosomal biogenesis in macrophages (By similarity). Also participates in immunologic synapse formation (PubMed:[27405666](#)).


Cellular Location Cell membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:P26041}; Cytoplasmic side {ECO:0000250|UniProtKB:P26041}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P26041}. Apical cell membrane {ECO:0000250|UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250|UniProtKB:P26041}; Cytoplasmic side {ECO:0000250|UniProtKB:P26041}. Cell projection, microvillus membrane {ECO:0000250|UniProtKB:P26041}; Peripheral membrane protein {ECO:0000250|UniProtKB:P26041}; Cytoplasmic side {ECO:0000250|UniProtKB:P26041}. Cell projection, microvillus {ECO:0000250|UniProtKB:P26041}. Note=Phosphorylated form is enriched in microvilli-like structures at apical membrane. Increased cell membrane localization of both phosphorylated and non-phosphorylated forms seen after thrombin treatment (By similarity). Localizes at the uropods of T lymphoblasts. {ECO:0000250|UniProtKB:P26041, ECO:0000269|PubMed:18586956, ECO:0000269|PubMed:9298994}

Tissue Location In all tissues and cultured cells studied.

Images



Western blot analysis of Moesin expression in HeLa cell lysate.



Immunohistochemical analysis of paraffin-embedded human thyroid cancer, using Moesin Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.