

Anti-ALDH1A1 Rabbit Monoclonal Antibody

Catalog # ABO13884

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P00352
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-ALDH1A1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	216
Other Names	Aldehyde dehydrogenase 1A1, 1.2.1.19, ALDH1A1 (HGNC:402)
Calculated MW	54862
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Subcellular Localization	Cytoplasm.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AHA-1
Immunogen	A synthesized peptide derived from human ALDH1A1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	ALDH1A1 (HGNC:402)
-------------	--------------------------------------

Function

Cytosolic dehydrogenase that catalyzes the irreversible oxidation of a wide range of aldehydes to their corresponding carboxylic acid (PubMed:[12941160](#), PubMed:[15623782](#), PubMed:[17175089](#), PubMed:[19296407](#), PubMed:[25450233](#), PubMed:[26373694](#), PubMed:[25413692](#)). Functions downstream of retinol dehydrogenases and catalyzes the oxidation of retinaldehyde into retinoic acid, the second step in the oxidation of retinol/vitamin A into retinoic acid (PubMed:[25413692](#)). This pathway is crucial to control the levels of retinol and retinoic acid, two important molecules which excess can be teratogenic and cytotoxic (PubMed:[25413692](#)). Also oxidizes aldehydes resulting from lipid peroxidation like (E)-4-hydroxynon-2-enal/HNE, malonaldehyde and hexanal that form protein adducts and are highly cytotoxic. By participating for instance to the clearance of (E)-4-hydroxynon-2-enal/HNE in the lens epithelium prevents the formation of HNE-protein adducts and lens opacification (PubMed:[12941160](#), PubMed:[15623782](#), PubMed:[19296407](#)). Also functions downstream of fructosamine-3-kinase in the fructosamine degradation pathway by catalyzing the oxidation of 3- deoxyglucosone, the carbohydrate product of fructosamine 3-phosphate decomposition, which is itself a potent glycating agent that may react with lysine and arginine side-chains of proteins (PubMed:[17175089](#)). Also has an aminobutyraldehyde dehydrogenase activity and is probably part of an alternative pathway for the biosynthesis of GABA/4-aminobutanoate in midbrain, thereby playing a role in GABAergic synaptic transmission (By similarity).


Cellular Location

Cytoplasm, cytosol. Cell projection, axon {ECO:0000250 | UniProtKB:P24549}


Tissue Location

Expressed by erythrocytes (at protein level).

Images



Western blot analysis of ALDH1A1 expression in HepG2 cell lysate.



Immunohistochemical analysis of paraffin-embedded human liver cancer, using ALDH1A1 Antibody.