

Anti-IL1 beta Rabbit Monoclonal Antibody

Catalog # ABO13918

Product Information

Application	WB
Primary Accession	P01584
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-IL1 beta Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.

Additional Information


Gene ID	3553
Other Names	Interleukin-1 beta, IL-1 beta, Catabolin, IL1B (HGNC:5992), IL1F2
Calculated MW	30748
Application Details	WB 1:500-1:2000
Subcellular Localization	Secreted. The lack of a specific hydrophobic segment in the precursor sequence suggests that IL-1 is released by damaged cells or is secreted by a mechanism differing from that used for other secretory proteins.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: BII-9
Immunogen	A synthesized peptide derived from human IL1 beta
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	IL1B (HGNC:5992)
Synonyms	IL1F2

Function	Potent pro-inflammatory cytokine (PubMed: 10653850 , PubMed: 12794819 , PubMed: 28331908 , PubMed: 3920526). Initially discovered as the major endogenous pyrogen, induces prostaglandin synthesis, neutrophil influx and activation, T-cell activation and cytokine production, B-cell activation and antibody production, and fibroblast proliferation and collagen production (PubMed: 3920526). Promotes Th17 differentiation of T-cells. Synergizes with IL12/interleukin-12 to induce IFNG synthesis from T-helper 1 (Th1) cells (PubMed: 10653850). Plays a role in angiogenesis by inducing VEGF production synergistically with TNF and IL6 (PubMed: 12794819). Involved in transduction of inflammation downstream of pyroptosis: its mature form is specifically released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed: 33377178 , PubMed: 33883744). Acts as a sensor of <i>S.pyogenes</i> infection in skin: cleaved and activated by pyogenes SpeB protease, leading to an inflammatory response that prevents bacterial growth during invasive skin infection (PubMed: 28331908).
Cellular Location	Cytoplasm, cytosol. Secreted. Lysosome Secreted, extracellular exosome {ECO:0000250 UniProtKB:P10749} Note=The precursor is cytosolic (PubMed:15192144). In response to inflammasome-activating signals, such as ATP for NLRP3 inflammasome or bacterial flagellin for NLRC4 inflammasome, cleaved and secreted (PubMed:24201029, PubMed:33377178, PubMed:33883744). Mature form is secreted and released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33883744). In contrast, the precursor form is not released, due to the presence of an acidic region that is proteolytically removed by CASP1 during maturation (PubMed:33883744). The secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10 (PubMed:32272059)
Tissue Location	Expressed in activated monocytes/macrophages (at protein level).

Images



Western blot analysis of extracts from recombinant IL1 beta, using IL1 beta antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.