

Anti-JNK1/2/3 MAPK8 Rabbit Monoclonal Antibody

Catalog # ABO13920

Product Information

Application	WB, IF, ICC, IP
Primary Accession	P45983/P45984/P53779
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-JNK1/2/3 MAPK8 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Calculated MW	48296 Da
Application Details	WB 1:1000-1:2000 ICC/IF 1:50-1:200 IP 1:50
Subcellular Localization	Cytoplasm. Nucleus.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5 mg/ml BSA.
Clone Names	Clone: DFI-13
Immunogen	A synthesized peptide derived from human JNK1+JNK2+JNK3
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Images

Immunofluorescent analysis using the Antibody at 1:50 dilution.



Immunofluorescent analysis of HeLa cells, using JNK1/2/3 Antibody.




Figure 1. Western blot analysis of JNK1/2/3 using anti-JNK1/2/3 antibody (M02608-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HeLa whole cell lysates,

Lane 2: human U87 whole cell lysates,

Lane 3: human A549 whole cell lysates,

Lane 4: rat PC-12 whole cell lysates,

Lane 5: rat C6 whole cell lysates,

Lane 6: mouse Neuro-2a whole cell lysates,

Lane 7: mouse NIH/3T3 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit

anti-JNK1/2/3 antigen affinity purified monoclonal antibody (Catalog # M02608-1) at 1:1000 overnight at 4°C,

then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP

secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced

Chemiluminescent detection (ECL) kit (Catalog # EK1002)

with Tanon 5200 system. A specific band was detected for JNK1/2/3 at approximately 40, 48 kDa. The expected band

size for JNK1/2/3 is at 48 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.