

Anti-Hepcidin HAMP Rabbit Monoclonal Antibody

Catalog # ABO13945

Product Information

Application	WB
Primary Accession	P81172
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Hepcidin HAMP Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human, Rat.


Additional Information

Gene ID	57817
Other Names	Hepcidin, Liver-expressed antimicrobial peptide 1, LEAP-1, Putative liver tumor regressor, PLTR, Hepcidin-25, Hepc25, Hepcidin-20, Hepc20, HAMP (HGNC:15598), HEPC, LEAP1
Calculated MW	9408
Application Details	WB 1:500-1:2000
Subcellular Localization	Secreted.
Tissue Specificity	Highest expression in liver and to a lesser extent in heart and brain. Low levels in lung, tonsils, salivary gland, trachea, prostate gland, adrenal gland and thyroid gland. Secreted into the urine..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AAHH-8
Immunogen	A synthesized peptide derived from human Hepcidin
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	HAMP (HGNC:15598)
Synonyms	HEPC, LEAP1
Function	Liver-produced hormone that constitutes the main circulating regulator of iron absorption and distribution across tissues. Acts by promoting endocytosis and degradation of ferroportin/SLC40A1, leading to the retention of iron in iron-exporting cells and decreased flow of iron into plasma (PubMed: 22682227 , PubMed: 29237594 , PubMed: 32814342). Controls the major flows of iron into plasma: absorption of dietary iron in the intestine, recycling of iron by macrophages, which phagocytose old erythrocytes and other cells, and mobilization of stored iron from hepatocytes (PubMed: 22306005).
Cellular Location	Secreted.
Tissue Location	Highest expression in liver and to a lesser extent in heart and brain. Low levels in lung, tonsils, salivary gland, trachea, prostate gland, adrenal gland and thyroid gland. Secreted into the urine and blood (PubMed:11034317). Expressed by hepatocytes (PubMed:15124018).

Images



Western blot analysis of Hepcidin expression in Human Hepcidin Full-length Recombinant Protein (GFP Tagged).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.