

Anti-HIF-1 beta ARNT Rabbit Monoclonal Antibody

Catalog # ABO14031

Product Information

Application	WB, IHC
Primary Accession	P27540
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-HIF-1 beta ARNT Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

Additional Information


Gene ID	405
Other Names	Aryl hydrocarbon receptor nuclear translocator, ARNT protein, Class E basic helix-loop-helix protein 2, bHLHe2, Dioxin receptor, nuclear translocator, Hypoxia-inducible factor 1-beta, HIF-1-beta, HIF1-beta, ARNT (HGNC:700), BHLHE2
Calculated MW	86636
Application Details	WB 1:500-1:2000 IHC 1:500-1:2000
Subcellular Localization	Nucleus.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: EAD-1
Immunogen	A synthesized peptide derived from human HIF-1 beta
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	ARNT (HGNC:700)
-------------	-----------------------------------

Synonyms	BHLHE2
Function	Transcription factor that functions as an obligate heterodimerization partner for several basic helix-loop-helix PER-ARNT- SIM (bHLH-PAS) proteins. Forms heterodimers with the aryl hydrocarbon receptor (AHR) to mediate the transcriptional response to xenobiotics by binding xenobiotic response elements (XREs) or dioxin response elements (DREs) in target gene promoters (PubMed: 28396409 , PubMed: 34521881). Also heterodimerizes with hypoxia-inducible factor alpha subunits (HIF1A, EPAS1, NPAS1 or NPAS3) to form hypoxia-inducible transcription factors that bind hypoxia response elements (HREs) and activate genes involved in cellular and systemic responses to hypoxia, including angiogenesis, erythropoiesis, and metabolic adaptation (PubMed: 16181639 , PubMed: 20699359). Plays a central role in environmental sensing and developmental and physiological regulation of gene expression (PubMed: 34521881).
Cellular Location	Nucleus.


Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of HIF-1 beta expression in Human fetal kidney lysate.



Immunohistochemical analysis of paraffin-embedded human colon, using HIF-1 beta Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.