

Anti-Aconitase 1 ACO1 Rabbit Monoclonal Antibody

Catalog # ABO14061

Product Information

Application	WB, IF, ICC
Primary Accession	P21399
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Aconitase 1 ACO1 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	48
Other Names	Cytoplasmic aconitate hydratase, Aconitase, 4.2.1.3, Citrate hydro-lyase, Ferritin repressor protein, Iron regulatory protein 1, IRP1, Iron-responsive element-binding protein 1, IRE-BP 1, ACO1, IREB1
Calculated MW	98399
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200
Subcellular Localization	Cytoplasm.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ABG-1
Immunogen	A synthesized peptide derived from human Aconitase 1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	ACO1
Synonyms	IREB1


Function

Bifunctional iron sensor that switches between 2 activities depending on iron availability (PubMed:[1281544](#), PubMed:[1946430](#), PubMed:[8041788](#)). Iron deprivation, promotes its mRNA binding activity through which it regulates the expression of genes involved in iron uptake, sequestration and utilization (PubMed:[1281544](#), PubMed:[1946430](#), PubMed:[23891004](#), PubMed:[8041788](#)). Binds to iron-responsive elements (IRES) in the untranslated region of target mRNAs preventing for instance the translation of ferritin and aminolevulinic acid synthase and stabilizing the transferrin receptor mRNA (PubMed:[1281544](#), PubMed:[1946430](#), PubMed:[23891004](#), PubMed:[8041788](#)).


Cellular Location

Cytoplasm, cytosol.


Images



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



Western blot analysis of Aconitase 1 in (1) mouse kidney lysate; (2) HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.