

Anti-GTPase HRAS Rabbit Monoclonal Antibody

Catalog # ABO14066

Product Information

Application	WB, IF, ICC, IP, FC
Primary Accession	P01112
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-GTPase HRAS Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Gene ID	3265
Other Names	GTPase HRas, 3.6.5.2, H-Ras-1, Ha-Ras, Transforming protein p21, c-H-ras, p21ras, GTPase HRas, N-terminally processed, HRAS, HRAS1
Calculated MW	21298
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50 FC 1:30
Subcellular Localization	Cell membrane. Cell membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus. Golgi apparatus membrane; Lipid-anchor. The active GTP-bound form is localized most strongly to membranes than the inactive GDP-bound form (By similarity). Shuttles between the plasma membrane and the Golgi apparatus..
Tissue Specificity	Widely expressed..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: BEA-8
Immunogen	A synthesized peptide derived from human GTPase HRAS
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	HRAS
Synonyms	HRAS1
Function	Signal transducer in the Ras-MAPK signaling pathway that regulates cell proliferation and survival (PubMed: 22821884). Ras proteins bind GDP/GTP and possess intrinsic GTPase activity (PubMed: 12740440 , PubMed: 14500341 , PubMed: 9020151). Recognized by LZTR1 that mediates its ubiquitination by a BCR (BTB-CUL3-RBX1) E3 ubiquitin- protein ligase complex (PubMed: 40934300).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P20171}; Lipid-anchor; Cytoplasmic side. Golgi apparatus. Golgi apparatus membrane; Lipid-anchor. Note=The active GTP-bound form is localized most strongly to membranes than the inactive GDP-bound form (By similarity). Shuttles between the plasma membrane and the Golgi apparatus.
Tissue Location	Widely expressed..

Images



Western blot analysis MCF7 cell lysate using HRAS Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.